

On projective representations for compact quantum groups

Kenny De Commer*

Dipartimento di Matematica, Università degli Studi di Roma Tor Vergata

Via della Ricerca Scientifica 1, 00133 Roma, Italy

e-mail: decommer@mat.uniroma2.it

Abstract

We study actions of compact quantum groups on type I factors, which may be interpreted as projective representations of compact quantum groups. We generalize to this setting some of Woronowicz's results concerning Peter-Weyl theory for compact quantum groups. The main new phenomenon is that for general compact quantum groups (more precisely, those which are not of Kac type), not all irreducible projective representations have to be finite-dimensional. As applications, we consider the theory of projective representations for the compact quantum groups associated with group von Neumann algebras of discrete groups, and consider a certain non-trivial projective representation for quantum $SU(2)$.

Keywords: compact quantum group; projective representation; Galois (co-)object

AMS 2010 *Mathematics subject classification:* 17B37, 81R50, 16T15

Introduction

It is well-known that for compact groups, one can easily extend the main theorems of the Peter-Weyl theory to cover also *projective* representations. In this article, we will see that if one tries to do the same for Woronowicz's compact *quantum* groups, one confronts at least one surprising novelty: not all irreducible projective representations of a compact quantum group have to be finite-dimensional. On the other hand, one will still be able to decompose any projective representation into a direct sum of irreducible ones, and to determine certain orthogonality relations between the matrix coefficients of irreducible projective representations.

The main tool we will use in this article are the *Galois co-objects* which we introduced in [10]. Indeed, we showed there that when one quantizes the notion of a projective representation, this structure plays the role of a 'generalized 2-cocycle function'.

In the *first* section of this article, we will develop a structure theory for such Galois co-objects in the setting of compact quantum groups. A lot of the techniques we use are directly inspired by the theory of the compact quantum groups themselves.

*Supported in part by the ERC Advanced Grant 227458 OACFT "Operator Algebras and Conformal Field Theory"

In the *second section*, we will show that such Galois co-objects can be dualized into Galois *objects* for their dual discrete quantum groups, a concept which was introduced in [8].

In the *third section*, which, except for the last part, is independent from the more technical second section, we present a Peter-Weyl theory for projective representations of compact quantum groups. We also show how projective representations give rise to module categories over the tensor category of the (ordinary) representations, and introduce the notion of *fusion rules* between irreducible projective representations and (ordinary) irreducible representations.

In the *fourth section*, we will give some details on the ‘reflection technique’ introduced in [10]. We showed there that from any Galois co-object for a given compact quantum group, one can create a (possibly) new *locally compact* quantum group. We will show that the type of this quantum group (namely whether it is compact or not) is intimately tied up with the behavior of the Galois co-object itself.

In the *fifth section*, we will consider the special case of compact *Kac* algebras. We show that in this case, all irreducible projective representations will be finite-dimensional, and the theory becomes essentially algebraic.

In the *sixth and seventh section*, we further specialize. We first quickly consider the case of *finite quantum groups* (i.e. *finite-dimensional Kac* algebras), for which we can mostly refer to the existing literature. Then we will treat *co-commutative* compact Kac algebras, which correspond to group von Neumann algebras of discrete groups. In this case, the projective representations turn out to be classified by certain special 2-cohomology classes of *finite subgroups* of the associated discrete group. In particular, we will be able to deduce that the group von Neumann algebra of a torsionless discrete group admits no non-trivial 2-cocycles. These results will be proven using only the material in the first section and the first part of the third section.

In the *eighth section*, we give a concrete example of what can happen in the non-Kac case by considering a particular non-trivial Galois co-object for the compact quantum group $SU_q(2)$. We compute explicitly all its associated projective representations, provide the corresponding orthogonality relations and calculate the fusion rules.

Notations and conventions

We will assume that all our Hilbert spaces are separable, and we take the inner product to be conjugate linear in the second argument. We also assume that all our von Neumann algebras have separable predual.

By ι , we denote the identity map on a set.

We denote by \odot the algebraic tensor product between vector spaces, by \otimes the tensor product between Hilbert spaces, and by $\bar{\otimes}$ the spatial tensor product between von Neumann algebras.

By Σ we denote the flip map between a tensor product of Hilbert spaces:

$$\Sigma : \mathcal{H}_1 \otimes \mathcal{H}_2 \rightarrow \mathcal{H}_2 \otimes \mathcal{H}_1 : \xi_1 \otimes \xi_2 \rightarrow \xi_2 \otimes \xi_1.$$

When $A \subseteq B(\mathcal{H}_1, \mathcal{H}_2)$ and $B \subseteq B(\mathcal{H}_2, \mathcal{H}_3)$ are linear spaces of maps between certain Hilbert spaces, we will denote $B \cdot A = \{\sum_{i=1}^n b_i a_i \mid n \in \mathbb{N}_0, b_i \in B, a_i \in A\}$.

We use the *leg numbering notation* for operators on tensor products of Hilbert spaces. E.g., if $Z : \mathcal{H}^{\otimes 2} \rightarrow \mathcal{H}^{\otimes 2}$ is a certain operator, we denote by Z_{13} the operator $\mathcal{H}^{\otimes 3} \rightarrow \mathcal{H}^{\otimes 3}$ acting as Z on the first and third factor, and as the identity on the second factor.

At certain points, we will need the theory of weights on von Neumann algebras, which is treated in detail in the first chapters of [27]. When M is a von Neumann algebra, and $\varphi : M^+ \rightarrow [0, \infty]$ is a normal semi-finite faithful (nsf) weight on M , we denote

$$\mathcal{N}_\varphi = \{x \in M \mid \varphi(x^*x) < \infty\}$$

for the space of square integrable elements, \mathcal{M}_φ^+ for the space of positive integrable elements, and \mathcal{M}_φ for the linear span of \mathcal{M}_φ^+ .

1 Galois co-objects for compact quantum groups

We begin with introducing the following concepts.

Definition 1.1. A von Neumann bialgebra (M, Δ_M) consists of a von Neumann algebra M and a faithful normal unital $*$ -homomorphism $\Delta_M : M \rightarrow M \bar{\otimes} M$, satisfying the coassociativity condition

$$(\Delta_M \otimes \iota)\Delta_M = (\iota \otimes \Delta_M)\Delta_M.$$

A von Neumann bialgebra (M, Δ_M) is called a compact Woronowicz algebra ([23],[20]) if there exists a normal state φ_M on M which is Δ_M -invariant:

$$(\varphi_M \otimes \iota)\Delta_M(x) = (\iota \otimes \varphi_M)\Delta_M(x) = \varphi_M(x)1 \quad \text{for all } x \in M.$$

A compact Woronowicz algebra is called a compact Kac algebra if there exists a normal Δ_M -invariant tracial state τ_M on M .

Remarks:

1. von Neumann bialgebras are also referred to as Hopf-von Neumann algebras in the literature. However, we prefer the above terminology, as for example a finite-dimensional Hopf-von Neumann algebra is not necessarily a Hopf algebra. Admittedly, a finite-dimensional von Neumann bialgebra is also not necessarily a bialgebra, as there could be no co-unit, but this seems a lesser ambiguity.
2. It is easy to see that a normal Δ_M -invariant state on a von Neumann bialgebra (M, Δ_M) , when it exists, is unique. One can moreover show that this state will automatically be faithful. We will then always use the notation φ_M for it in the general setting, but use the notation τ_M in the setting of compact Kac algebras to emphasize the traciality.
3. Compact Woronowicz algebras can be characterized as those von Neumann bialgebras arising from Woronowicz's compact quantum groups in the C^* -algebra setting ([34],[36]), by performing a GNS-type construction. However, we have decided to focus only on the von Neumann algebraic picture in this paper.

Let us also introduce the following notations, which will be constantly used in the following.

Notation 1.2. Let (M, Δ_M) be a compact Woronowicz algebra. We denote by $(\mathcal{L}^2(M), \cdot, \Delta_M)$ the GNS-construction of M with respect to φ_M . That is, $\mathcal{L}^2(M)$ is the completion of M , considered as a pre-Hilbert space with respect to the inner product structure

$$\langle x, y \rangle = \varphi_M(y^*x),$$

and Λ_M is the natural inclusion $M \hookrightarrow \mathcal{L}^2(M)$. We then identify M as a von Neumann subalgebra of $B(\mathcal{L}^2(M))$ by letting $x \in M$ corresponding to the (bounded) closure of the operator

$$\Lambda_M(M) \rightarrow \mathcal{L}^2(M) : \Lambda_M(y) \rightarrow \Lambda_M(xy), \quad \text{for all } y \in M.$$

We will further denote by ξ_M the cyclic and separating vector $\Lambda_M(1_M)$ in $\mathcal{L}^2(M)$, so that $x\xi_M = \Lambda_M(x)$ for all $x \in M$.

The following two unitaries are of fundamental importance.

Definition 1.3. Let (M, Δ_M) be a compact Woronowicz algebra.

The right regular corepresentation of (M, Δ_M) is defined to be the unitary $V \in B(\mathcal{L}^2(M)) \bar{\otimes} M$ which is uniquely determined by the formula

$$V\Lambda_M(x) \otimes \eta = \Delta_M(x)\xi_M \otimes \eta, \quad \text{for all } x \in M, \eta \in \mathcal{L}^2(M).$$

The left regular corepresentation of (M, Δ_M) is defined to be the unitary $W \in M \bar{\otimes} B(\mathcal{L}^2(M))$ which is uniquely determined by the fact that

$$W^*\eta \otimes \Lambda_M(x) = \Delta_M(x)\eta \otimes \xi_M, \quad \text{for all } x \in M, \eta \in \mathcal{L}^2(M).$$

We will in the following always use the above notations for these corepresentations. Note that establishing the unitarity of these maps requires some non-trivial work! An approach to compact quantum groups based on the properties of such unitaries can be found in [3], section 4.

Let us now introduce the notion of a Galois co-object for a compact Woronowicz algebra (see [10]).

Definition 1.4. Let (M, Δ_M) be a compact Woronowicz algebra. A right Galois co-object for (M, Δ_M) consists of a Hilbert space $\mathcal{L}^2(N)$, a σ -weakly closed linear space $N \subseteq B(\mathcal{L}^2(M), \mathcal{L}^2(N))$ and a normal linear map $\Delta_N : N \rightarrow N \bar{\otimes} N$, such that the following properties hold: with N^{op} denoting the set

$$N^{op} := \{x^* \mid x \in N\} \subseteq B(\mathcal{L}^2(N), \mathcal{L}^2(M)),$$

we should have

1. $N \cdot \mathcal{L}^2(M)$ is norm-dense in $\mathcal{L}^2(N)$, and $N^{op} \cdot \mathcal{L}^2(N)$ is norm-dense in $\mathcal{L}^2(M)$,
2. the space N is a right M -module (by composition of operators),
3. for each $x, y \in N$, we have $x^*y \in M$,
4. $\Delta_N(xy) = \Delta_N(x)\Delta_M(y)$ for all $x \in N$ and $y \in M$,
5. $\Delta_N(x)^*\Delta_N(y) = \Delta_M(x^*y)$ for all $x, y \in N$,
6. Δ_N is coassociative: $(\Delta_N \otimes \iota)\Delta_N = (\iota \otimes \Delta_N)\Delta_N$, and
7. the linear span of the set $\{\Delta_N(x)(y \otimes z) \mid x \in N, y, z \in M\}$ is σ -weakly dense in $N \bar{\otimes} N$.

If (N_1, Δ_{N_1}) and (N_2, Δ_{N_2}) are two Galois co-objects for a von Neumann bialgebra (M, Δ_M) , we call them isomorphic if there exists a unitary $u : \mathcal{L}^2(N_1) \rightarrow \mathcal{L}^2(N_2)$ such that $uN_1 = N_2$ and

$$\Delta_{N_2}(ux) = (u \otimes u)\Delta_{N_1}(x) \quad \text{for all } x \in N_1.$$

Remarks:

1. The previous definition can be shown to be equivalent with the one presented in [10], Definition 0.5. Also remark that the previous conditions can be grouped together as follows: a Galois co-object is a right Morita (or imprimitivity) Hilbert M -module (conditions 1 to 3) with a Δ_M -compatible coalgebra structure (conditions 4 and 5 and condition 6) which is ‘non-degenerate’ (condition 7).

2. A trivial example of a right (M, Δ_M) -Galois co-object is (M, Δ_M) itself. Indeed, the final condition even holds in a stronger form, as it can be shown that already $\{\Delta_M(x)(1 \otimes y) \mid x, y \in M\}$ is σ -weakly dense in $M \bar{\otimes} M$ for compact Woronowicz algebras. It follows that this stronger condition is then in fact true for *all* Galois co-objects for compact Woronowicz algebras.

3. A treatment of Galois co-objects in the setting of Hopf algebras can be found in [6].

One can similarly define the notion of a *left* Galois co-object. Left Galois co-objects can be created from right ones in the following way.

Definition 1.5. *Let (N, Δ_N) be a right Galois co-object for the compact Woronowicz algebra (M, Δ_M) .*

We call the couple $(N^{op}, \Delta_{N^{op}})$, consisting of

$$N^{op} = \{x^* \mid x \in N\} \subseteq B(\mathcal{L}^2(N), \mathcal{L}^2(M)),$$

together with the coproduct

$$\Delta_{N^{op}}(x) := \Delta_N(x^*)^*, \quad x \in N^{op},$$

the opposite (left) Galois co-object of (N, Δ_N) . It is a left Galois co-object for the compact Woronowicz algebra (M, Δ_M) .

We call the couple $(N^{cop}, \Delta_{N^{cop}})$, where

$$N^{cop} = N \subseteq B(\mathcal{L}^2(M), \mathcal{L}^2(N))$$

and

$$\Delta_{N^{cop}} = \Delta_N^{op} : N \rightarrow N \bar{\otimes} N : x \rightarrow \Sigma \Delta_N(x) \Sigma, \quad \text{for all } x \in N,$$

the co-opposite (right) Galois co-object of (N, Δ_N) . It is a right Galois co-object for the compact Woronowicz algebra (M, Δ_M^{op}) .

The following notation will be useful.

Notation 1.6. *Let (M, Δ_M) be a compact Woronowicz algebra, and (N, Δ_N) a right Galois co-object for (M, Δ_M) . We denote*

$$\Lambda_N : N \rightarrow \mathcal{L}^2(N) : x \rightarrow x \xi_M.$$

Remark: By the second condition in Definition 1.4, we know that N is a right M -module, and then we trivially have that

$$\Lambda_N(xy) = x \Lambda_M(y), \quad \text{for all } x \in N, y \in M.$$

By the third condition in that definition, together with the faithfulness of φ_M , we see that Λ_N is injective and that

$$\langle \Lambda_N(x), \Lambda_N(y) \rangle = \varphi_M(y^* x), \quad \text{for all } x, y \in N.$$

And finally, by the first (and second) condition in that definition, we see that Λ_N has norm-dense range.

One can construct for a Galois co-object (N, Δ_N) certain unitaries which are analogous to the regular corepresentations for a compact Woronowicz algebra (and coincide with them in case $(N, \Delta_N) = (M, \Delta_M)$).

Proposition 1.7. *Let (M, Δ_M) be a compact Woronowicz algebra, and (N, Δ_N) a right Galois co-object for (M, Δ_M) .*

1. There exists a unitary

$$\tilde{V} : \mathcal{L}^2(N) \otimes \mathcal{L}^2(M) \rightarrow \mathcal{L}^2(N) \otimes \mathcal{L}^2(N)$$

which is uniquely determined by the property that for all $\eta \in \mathcal{L}^2(M)$ and $x \in N$, we have

$$\tilde{V} \Lambda_N(x) \otimes \eta = \Delta_N(x) \xi_M \otimes \eta.$$

Similarly, there exists a unitary

$$\tilde{W} : \mathcal{L}^2(N) \otimes \mathcal{L}^2(N) \rightarrow \mathcal{L}^2(M) \otimes \mathcal{L}^2(N),$$

uniquely determined by the property that for all $\eta \in \mathcal{L}^2(M)$ and $x \in N$, we have

$$\tilde{W}^* \eta \otimes \Lambda_N(x) = \Delta_N(x) \eta \otimes \xi_M.$$

2. We have $\tilde{V} \in B(\mathcal{L}^2(N)) \bar{\otimes} N$ and $\tilde{W}^* \in N \bar{\otimes} B(\mathcal{L}^2(N))$.

3. For $x \in N$, we have

$$\Delta_N(x) = \tilde{V}(x \otimes 1)V^* = \tilde{W}^*(1 \otimes x)W.$$

4. The following ‘pentagonal identities’ hold:

$$\tilde{V}_{12} \tilde{V}_{13} V_{23} = \tilde{V}_{23} \tilde{V}_{12}$$

as maps from $\mathcal{L}^2(N) \otimes \mathcal{L}^2(M) \otimes \mathcal{L}^2(M)$ to $\mathcal{L}^2(N) \otimes \mathcal{L}^2(N) \otimes \mathcal{L}^2(N)$, and

$$W_{12} \tilde{W}_{13} \tilde{W}_{23} = \tilde{W}_{23} \tilde{W}_{12}$$

as maps from $\mathcal{L}^2(N) \otimes \mathcal{L}^2(N) \otimes \mathcal{L}^2(N)$ to $\mathcal{L}^2(M) \otimes \mathcal{L}^2(M) \otimes \mathcal{L}^2(N)$.

5. The following identities hold:

$$\begin{aligned} (\iota \otimes \Delta_N) \tilde{V} &= \tilde{V}_{12} \tilde{V}_{13}, \\ (\Delta_N \otimes \iota)(\tilde{W}^*) &= \tilde{W}_{23}^* \tilde{W}_{13}^*. \end{aligned}$$

Proof. The statements for \tilde{W} follow immediately from the ones for \tilde{V} , by considering the co-opposite Galois co-object.

We then refer to [10] for the proofs of the first four statements (Proposition 2.3 for the first and second assertion, Proposition 2.4 for the third and fourth). The fifth statement follows immediately from combining the three preceding ones. \square

Remark: Although we referred to [10], we want to stress that these assertions are quite straightforward to prove. For example, the surjectivity of \tilde{V} follows quite immediately from the seventh condition in Definition 1.4, combined with the surjectivity of V .

Definition 1.8. We call the unitary \tilde{V} appearing in the previous proposition the right regular (N, Δ_N) -corepresentation of (M, Δ_M) . We call the unitary \tilde{W} the left regular $(N^{\text{op}}, \Delta_{N^{\text{op}}})$ -corepresentation of (M, Δ_M) (where we recall that $(N^{\text{op}}, \Delta_{N^{\text{op}}})$ is the left Galois co-object opposite to (N, Δ_N) , see Definition 1.5).

Remark: The general notion of an ‘ (N, Δ_N) -corepresentation’ will be introduced in the third section.

For the rest of this section, we will fix a compact Woronowicz algebra (M, Δ_M) and a right Galois co-object (N, Δ_N) for (M, Δ_M) . We then further keep denoting by V and W the right and left regular corepresentations of (M, Δ_M) , and by \tilde{V} and \tilde{W} the right regular (N, Δ_N) - and left regular $(N^{\text{op}}, \Delta_{N^{\text{op}}})$ -corepresentation of (M, Δ_M) .

Our following lemma improves the second assertion in Proposition 1.7.

Lemma 1.9. *The following equalities hold:*

$$\begin{aligned} N &= \{(\omega \otimes \iota)(\tilde{V}) \mid \omega \in B(\mathcal{L}^2(N))_*\}^{\sigma\text{-weak closure}} \\ &= \{(\iota \otimes \omega)(\tilde{W}^*) \mid \omega \in B(\mathcal{L}^2(N))_*\}^{\sigma\text{-weak closure}}. \end{aligned}$$

Proof. We will again only prove the first identity, as the second one then follows by symmetry.

For $\xi, \eta \in \mathcal{L}^2(N)$, denote by $\omega_{\xi, \eta}$ the normal functional on $B(\mathcal{L}^2(N))$ determined by $\omega_{\xi, \eta}(x) = \langle x\xi, \eta \rangle$ for $x \in B(\mathcal{L}^2(N))$. Then for $x, y \in N$, a straightforward computation shows that

$$(\omega_{\Delta_N(x), \Delta_N(y)} \otimes \iota)(\tilde{V}) = (\varphi_M \otimes \iota)((y^* \otimes 1)\Delta_N(x)).$$

It is thus enough to prove that the linear span of such elements is σ -weakly dense in N .

Suppose that this were not so. Then we could find a non-zero $\omega \in N_*$ such that

$$\varphi_M(y^*(\iota \otimes \omega)(\Delta_N(x))) = 0 \quad \text{for all } x, y \in N.$$

Taking y equal to $(\iota \otimes \omega)(\Delta_N(x))$, we would have $(\iota \otimes \omega)(\Delta_N(x)) = 0$ for all $x \in N$ by faithfulness of φ_M . But then also

$$(\iota \otimes \omega)(\Delta_N(x)(m \otimes 1)) = 0 \quad \text{for all } x \in N, m \in M.$$

Now the set $\{\Delta_M(m_1)(m_2 \otimes 1) \mid m_1, m_2 \in M\}$ has σ -weakly dense linear span in $M\bar{\otimes}M$. It then follows, by the conditions 2, 4 and 7 in Definition 1.4 that

$$(\iota \otimes \omega)(z) = 0 \quad \text{for all } z \in N\bar{\otimes}N,$$

and so necessarily $\omega = 0$, a contradiction. □

The following result will allow us to obtain a decomposition for \tilde{W} and \tilde{V} .

Proposition 1.10. *Denote by $\hat{N} \subseteq B(\mathcal{L}^2(N))$ the von Neumann algebra*

$$\hat{N} = \{x \in B(\mathcal{L}^2(N)) \mid \tilde{V}^*(x \otimes 1)\tilde{V} = x \otimes 1\}.$$

Then \hat{N} satisfies the following properties.

1. *The von Neumann algebra \hat{N} is an l^∞ -sum of type I-factors.*
2. *The equality $\hat{N} = \{(\omega \otimes \iota)(\tilde{W}^*) \mid \omega \in B(\mathcal{L}^2(M), \mathcal{L}^2(N))_*\}^{\sigma\text{-weak closure}}$ holds.*

Remark: In the special case where (N, Δ_N) equals (M, Δ_M) considered as a right Galois co-object over itself, one denotes the above von Neumann algebra as \widehat{M} .

Proof. Consider the unital normal faithful $*$ -homomorphism

$$\text{Ad}_L : B(\mathcal{L}^2(N)) \rightarrow M \bar{\otimes} B(\mathcal{L}^2(N)) : x \rightarrow \Sigma \tilde{V}^*(x \otimes 1) \tilde{V} \Sigma.$$

Then by Proposition 1.7.5, it follows that Ad_L is a coaction by (M, Δ_M) :

$$(\Delta_M \otimes \iota) \text{Ad}_L = (\iota \otimes \text{Ad}_L) \text{Ad}_L.$$

Hence \hat{N} is precisely the set $B(\mathcal{L}^2(N))^{\text{Ad}_L}$ of Ad_L -fixed elements in $B(\mathcal{L}^2(N))$, that is, the set of elements satisfying $\text{Ad}_L(x) = 1 \otimes x$. It is well-known (and easy to see) that the map

$$E : B(\mathcal{L}^2(N)) \rightarrow B(\mathcal{L}^2(N)) : x \rightarrow (\varphi_M \otimes \iota) \text{Ad}_L(x)$$

is then a *normal* conditional expectation of $B(\mathcal{L}^2(N))$ onto \hat{N} . This forces \hat{N} to be an l^∞ -direct sum of type I -factors (see for example Exercise IX.4.1 in [27]).

We now prove the second point. First of all, remark that

$$\tilde{V}_{23} \tilde{W}_{12}^* = \tilde{W}_{12}^* \tilde{V}_{23},$$

which follows from a straightforward computation. From this, it is easy to get that

$$(\iota \otimes \text{Ad}_L)(\tilde{W}) = \tilde{W}_{13},$$

and so all elements of the form $(\omega \otimes \iota)(\tilde{W})$ with $\omega \in B(\mathcal{L}^2(N), \mathcal{L}^2(M))_*$ lie in \hat{N} . We next show that all elements of \hat{N} can be approximated σ -weakly by such elements.

For $\omega_1, \omega_2 \in N_*$, denote

$$\omega_1 * \omega_2 := (\omega_1 \otimes \omega_2) \circ \Delta_N \in N_*.$$

For $\xi, \eta \in B(\mathcal{L}^2(N))$, denote $\theta_{\xi, \eta}$ for the rank one operator $\zeta \rightarrow \langle \zeta, \eta \rangle \xi$ on $\mathcal{L}^2(N)$, and denote $\omega_{\xi, \eta}$ for the normal functional $x \rightarrow \langle x \xi, \eta \rangle$. Choose $b, x, y \in N$, and denote

$$a = (\omega_{\Lambda_N(x), \Lambda_N(y)} \otimes \iota)(\tilde{V}) \in N, \quad S_N(a) = (\omega_{\Lambda_N(x), \Lambda_N(y)} \otimes \iota)(\tilde{V}^*) \in N^{\text{op}},$$

where we recall that $N^{\text{op}} = \{x^* \mid x \in N\} \subseteq B(\mathcal{L}^2(N), \mathcal{L}^2(M))$. We will prove the identity

$$E(\theta_{\Lambda_N(a), \Lambda_N(b)}) = ((\varphi_M(b^* \cdot) * \varphi_M(S_N(a) \cdot)) \otimes \iota)(\tilde{W}^*), \quad (1)$$

where E is the conditional expectation defined in the first part of the proof, and where $\varphi_M(b^* \cdot)$ and $\varphi_M(S_N(a) \cdot)$ are the obvious normal functionals on N . As the linear span of the $\theta_{\Lambda_N(a), \Lambda_N(b)}$ is σ -weakly dense in $B(\mathcal{L}^2(N))$ by Lemma 1.9, and as E is a normal map with \hat{N} as its range, the second point of the proposition will follow from this identity.

To prove the identity (1), choose further $c, d \in N$. It is sufficient to prove then that

$$\langle E(\theta_{\Lambda_N(a), \Lambda_N(b)}) \cdot \Lambda_N(c), \Lambda_N(d) \rangle = \langle ((\varphi_M(b^* \cdot) * \varphi_M(S_N(a) \cdot)) \otimes \iota)(\tilde{W}^*) \cdot \Lambda_N(c), \Lambda_N(d) \rangle. \quad (2)$$

We remark now that a and $S_N(a)$ can also be rewritten in the following form, by a simple computation involving only the definition of \tilde{V} :

$$a = (\varphi_N \otimes \iota)((y^* \otimes 1) \Delta_N(x)), \quad S_N(a) = (\varphi_M \otimes \iota)(\Delta_N(y)^*(x \otimes 1)).$$

Using again the definition of \tilde{V} , the left hand of equation (2) then simplifies to

$$(\varphi_M \otimes \varphi_M \otimes \varphi_M \otimes \varphi_M)((y^* \otimes 1 \otimes b^* \otimes 1) \Delta_N(d)_{24}^* \Delta_N(c)_{34} \Delta_N(x)_{12}). \quad (3)$$

On the other hand, using the definition of \widetilde{W}^* , we get that the right hand side of equation (2) becomes

$$(\varphi_M \otimes \varphi_M \otimes \varphi_M \otimes \varphi_M)((1 \otimes b^* \otimes 1 \otimes d^*)\Delta_N(y)_{13}^*(x \otimes \Delta_N^{(2)}(c))), \quad (4)$$

where $\Delta_N^{(2)}(c) = (\iota \otimes \Delta_N)\Delta_N(c)$. In both expressions (3) and (4), we can write

$$v = (\varphi_M \otimes \iota)((b^* \otimes 1)\Delta_N(c)),$$

and we then have to prove that

$$(\varphi_M \otimes \varphi_M \otimes \varphi_M)((y \otimes \Delta_N(d))^*(\Delta_N(x) \otimes v)) = (\varphi_M \otimes \varphi_M \otimes \varphi_M)((\Delta_N(y) \otimes d)^*(x \otimes \Delta_N(v))). \quad (5)$$

Now by the final condition in Definition 1.4 (and the second remark following it), it is enough to show that these two expressions are equal when we replace $x \otimes v$ by $\Delta_N(z)(m \otimes 1)$ and $y \otimes d$ by $\Delta_N(w)(n \otimes 1)$, where $w, z \in N$ and $m, n \in M$. But then the left hand side of (5) becomes

$$(\varphi_M \otimes \varphi_M \otimes \varphi_M)((n^* \otimes 1 \otimes 1)\Delta_M^{(2)}(w^*z)(\Delta_M(m) \otimes 1)),$$

which by invariance of φ_M collapses to $\varphi_M(n^*w^*zm)$. A similar computation shows that with this replacement, also the right hand side expression in (5) collapses to $\varphi_M(n^*w^*zm)$. This concludes the proof. \square

Of course, we then also have

$$\widehat{N} = \{(\omega \otimes \iota)(\widetilde{W}) \mid \omega \in B(\mathcal{L}^2(N), \mathcal{L}^2(M))_*\}^{\sigma\text{-weak closure}},$$

which follows immediately by applying the $*$ -operation to both sides of the identity in the second point of the previous proposition.

Notation 1.11. By Proposition 1.10.1, we may identify the center $\mathcal{Z}(\widehat{N})$ of \widehat{N} with $l^\infty(I_N)$, for some countable set I_N . Denoting p_r the minimal central projection in $\mathcal{Z}(\widehat{N})$ associated with the element $r \in I_N$, we may further identify $p_r\widehat{N}$ with $B(\mathcal{H}_r)$ for some separable Hilbert space \mathcal{H}_r . We also denote

$$n_r := \dim(\mathcal{H}_r) \in \mathbb{N}_0 \cup \{\infty\}.$$

Proposition 1.12. *The unital normal faithful $*$ -homomorphism*

$$\text{Ad}_R : \widehat{N} \rightarrow B(\mathcal{L}^2(N)) \overline{\otimes} M : x \rightarrow \Sigma \widetilde{W}(1 \otimes x) \widetilde{W}^* \Sigma$$

restricts to a $$ -homomorphism $\widehat{N} \rightarrow \widehat{N} \overline{\otimes} M$, and defines in this way a right coaction of (M, Δ_M) on \widehat{N} .*

Moreover, the set of fixed elements for Ad_R coincides with the center $\mathcal{Z}(\widehat{N})$ of \widehat{N} .

Proof. From Proposition 1.10.2, it follows that $\widetilde{W} \in N^{\text{op}} \overline{\otimes} \widehat{N}$ (we may apply the weak slice map property as N^{op} is a corner of a von Neumann algebra). Hence $\text{Ad}_R(x) \in \widehat{N} \overline{\otimes} M$ for $x \in \widehat{N}$. By applying Proposition 1.7.5, we get

$$(\text{Ad}_R \otimes \iota)\text{Ad}_R(x) = (\iota \otimes \Delta_M)\text{Ad}_R(x).$$

Hence the first part of the proposition follows.

If further $x \in \widehat{N}$ is a fixed element for Ad_R , then it follows that $(1 \otimes x)\widetilde{W} = \widetilde{W}(1 \otimes x)$. Again by Proposition 1.10.2, we deduce that $xy = yx$ for all $y \in \widehat{N}$, i.e. $x \in \mathcal{Z}(\widehat{N})$. \square

Corollary 1.13. *Using Notation 1.11 and the notation from the previous proposition, the coaction Ad_R restricts to an ergodic coaction*

$$Ad_R^{(r)} : B(\mathcal{H}_r) \rightarrow B(\mathcal{H}_r) \bar{\otimes} M$$

for each $r \in I_N$.

We recall that a coaction α is called ergodic if the only elements satisfying $\alpha(x) = x \otimes 1$ are scalar multiples of the unit element.

Proof. Clearly, as $\mathcal{Z}(\hat{N})$ consists of the fixed points of Ad_R by the previous proposition, it is immediate that Ad_R indeed restricts to $B(\mathcal{H}_r)$. If then x is a fixed element for $Ad_R^{(r)}$, we have, again by the previous proposition, that $x \in \mathcal{Z}(\hat{N}) \cap B(\mathcal{H}_r)$, and x is a scalar operator. \square

Now as each $Ad_R^{(r)}$ appearing in the previous corollary is ergodic, there exists a *unique* $Ad_R^{(r)}$ -invariant state $\phi_{N,r}$ on $B(\mathcal{H}_r)$, determined by the formula

$$\phi_{N,r}(x)1_{B(\mathcal{H}_r)} = (\iota \otimes \varphi_M)(Ad_R^{(r)}(x)), \quad \text{for all } x \in B(\mathcal{H}_r).$$

Notation 1.14. *If T_r is the positive trace class operator associated with the state $\phi_{N,r}$ on $B(\mathcal{H}_r)$ introduced above, we denote by $T_{r,0} \geq T_{r,1} \geq \dots$ the descending sequence of its eigenvalues, counting multiplicities. We further fix in \mathcal{H}_r a basis $e_{r,i}$, with $0 \leq i < n_r$, such that $e_{r,i}$ is an eigenvector for T_r with eigenvalue $T_{r,i}$.*

We denote by $e_{r,ij} \in \hat{N}$ the matrix units associated with the basis $e_{r,i}$, and we denote by $\omega_{r,ij}$ the following normal functionals on $\hat{N} \subseteq B(\bigoplus_{r \in I_N} \mathcal{H}_r)$:

$$\omega_{r,ij}(x) = \langle x e_{r,i}, e_{r,j} \rangle, \quad x \in \hat{N}.$$

In the special case where (N, Δ_N) equals (M, Δ_M) considered as a right Galois co-object over itself, we will denote the n_r as m_r , the $T_{r,j}$ as $D_{r,j}$ and the \mathcal{H}_r as \mathcal{K}_r , but otherwise keep all notation as above.

Theorem 1.15. *Denote*

$$\widetilde{W}_{r,ij} = (\iota \otimes \omega_{r,ji})(\widetilde{W}) \in N^{op} \subseteq B(\mathcal{L}^2(N), \mathcal{L}^2(M)).$$

Then the following statements hold.

1. The unitary \widetilde{W} equals the strong* convergent sum $\sum_{r \in I_N} \sum_{i,j=0}^{n_r-1} \widetilde{W}_{r,ij} \otimes e_{r,ij}$.
2. For each $r \in I_N$ and $0 \leq i, j < n_r$, we have

$$\sum_{k=0}^{n_r-1} \widetilde{W}_{r,ik} \cdot \widetilde{W}_{r,jk}^* = \delta_{i,j} 1_{\mathcal{L}^2(M)},$$

$$\sum_{k=0}^{n_r-1} \widetilde{W}_{r,ki}^* \cdot \widetilde{W}_{r,kj} = \delta_{i,j} 1_{\mathcal{L}^2(N)},$$

both sums converging strongly.

3. For each $r \in I_N$ and $0 \leq i, j < n_r$, we have

$$\Delta_N(\widetilde{W}_{r,ij}^*) = \sum_{k=0}^{n_r-1} \widetilde{W}_{r,ik}^* \otimes \widetilde{W}_{r,kj}^*,$$

the sum again being a strongly* converging one.

4. The following orthogonality relations hold:

$$\varphi_M(\widetilde{W}_{r,ij} \cdot \widetilde{W}_{s,kl}^*) = \delta_{r,s} \delta_{i,k} \delta_{j,l} T_{r,j}, \quad \text{for all } r, s \in I_N, 0 \leq i, j < n_r, 0 \leq k, l < n_s.$$

Proof. The first point is immediate, and also the second one follows straightforwardly from the unitality of \widetilde{W} . The third point follows from the identity $(\Delta_N \otimes \iota)(\widetilde{W}^*) = \widetilde{W}_{23}^* \widetilde{W}_{13}^*$ in Proposition 1.7.5. In the fourth point, the orthogonality relations for $r = s$ follow from writing out the identity

$$(\iota \otimes \varphi_M)(\text{Ad}_R^{(r)}(e_{r,ij})) = \phi_{N,r}(e_{r,ij}) = \delta_{i,j} T_{r,j}.$$

Thus the only thing left to show is that $\varphi_M(\widetilde{W}_{r,ij} \cdot \widetilde{W}_{s,kl}^*) = 0$ for $r \neq s$. But also here, we can use a standard technique (see e.g. [34]). For suppose that this were not so, and choose $r \neq s$ which violate this condition. Consider, for $x \in B(\mathcal{H}_s, \mathcal{H}_r)$, the element

$$F(x) = (\varphi_M \otimes \iota)(\widetilde{W}_r(1 \otimes x)\widetilde{W}_s^*) \in B(\mathcal{H}_s, \mathcal{H}_r),$$

where of course

$$\widetilde{W}_r = (1 \otimes p_r)\widetilde{W} = \sum_{i,j=0}^{n_r-1} \widetilde{W}_{r,ij} \otimes e_{r,ij} \in N^{\text{op}} \overline{\otimes} B(\mathcal{H}_r).$$

By assumption, there must exist an x with $F(x) \neq 0$. Fixing such an x , denote $y = F(x)$. Then it is easy to see that

$$\widetilde{W}_r(1 \otimes y)\widetilde{W}_s^* = (1 \otimes y),$$

using Proposition 1.7.5 and the Δ_M -invariance of φ_M . This implies that y^*y , resp. yy^* , is a fixed element for $\text{Ad}_R^{(s)}$, resp. $\text{Ad}_R^{(r)}$. Since these coactions are ergodic, y^*y and yy^* must be (identical) scalars, and so we can scale x such that y becomes a unitary u .

We then find that

$$(1 \otimes u)\widetilde{W}_s(1 \otimes u^*) = \widetilde{W}_r.$$

This implies that there exist two non-equal normal functionals ω_1 and ω_2 on \widehat{N} such that

$$(\iota \otimes \omega_1)(\widetilde{W}) = (\iota \otimes \omega_2)(\widetilde{W}).$$

As the set $\{(\omega \otimes \iota)(\widetilde{W}) \mid \omega \in B(\mathcal{L}^2(N), \mathcal{L}^2(M))\}$ is σ -weakly dense in \widehat{N} by Proposition 1.10.2, this clearly gives a contradiction. Hence $\varphi_M(\widetilde{W}_{r,ij} \cdot \widetilde{W}_{s,kl}^*) = 0$ for $r \neq s$. □

Notation 1.16. By the final part of the previous proposition, we have a unitary transformation

$$\mathcal{L}^2(N) \cong \bigoplus_{r \in I_N} \mathcal{H}_r \otimes \overline{\mathcal{H}_r},$$

by means of the map

$$\widetilde{W}_{r,ij}^* \xi_M \rightarrow T_{r,j}^{1/2} e_{r,i} \otimes \overline{e_{r,j}}.$$

In the following, we will then always identify $\mathcal{L}^2(N)$ and $\bigoplus_{r \in I_N} \mathcal{H}_r \otimes \overline{\mathcal{H}_r}$ in this way, so that for example the elements $x \in N$ act directly as linear operators

$$\bigoplus_{r \in I_M} \mathcal{H}_r \otimes \overline{\mathcal{H}_r} \rightarrow \bigoplus_{r \in I_N} \mathcal{H}_r \otimes \overline{\mathcal{H}_r}.$$

Lemma 1.17. 1. With $\tilde{V}_{r,ij} := T_{r,i}^{1/2} T_{r,j}^{-1/2} \tilde{W}_{r,ij}^*$, we have the identity

$$\tilde{V} = \sum_{r \in I_N} \sum_{i,j=0}^{n_r-1} \overline{e_{r,ij}} \otimes \tilde{V}_{r,ij},$$

the sum converging strongly*.

2. The $\tilde{V}_{r,ij}$ satisfy the following orthogonality relations:

$$\varphi_M(\tilde{V}_{r,ij}^* \tilde{V}_{s,kl}) = \delta_{r,s} \delta_{i,k} \delta_{j,l} T_{r,i} \quad \text{for all } r, s \in I_N, 0 \leq i, j < n_r, 0 \leq k, l < n_s.$$

3. The following equalities hold:

$$\begin{aligned} \hat{N}' &= \{(\iota \otimes \omega)(\tilde{V}) \mid \omega \in N_*\} \\ &= \{x \in B(\mathcal{L}^2(N)) \mid \tilde{W}^*(1 \otimes x)\tilde{W} = 1 \otimes x\}. \end{aligned}$$

Proof. Choose $r \in I_N$, $0 \leq i, j < n_r$ and $\eta \in \mathcal{L}^2(M)$. Then we compute

$$\begin{aligned} \tilde{V} e_{r,i} \otimes \overline{e_{r,j}} \otimes \eta &= T_{r,j}^{-1/2} \tilde{V}(\tilde{W}_{r,ij}^* \xi_M \otimes \eta) \\ &= T_{r,j}^{-1/2} \sum_{k=0}^{n_r-1} \tilde{W}_{r,ik}^* \xi_M \otimes \tilde{W}_{r,kj}^* \eta \\ &= \sum_{k=0}^{n_r-1} T_{r,k}^{1/2} T_{r,j}^{-1/2} e_{r,i} \otimes \overline{e_{r,k}} \otimes \tilde{W}_{r,kj}^* \eta. \end{aligned}$$

From this, the first point in the lemma follows.

The second point is of course just a reformulation of Theorem 1.15.4.

These orthogonality relations then immediately imply that

$$\hat{N}' = \{(\iota \otimes \omega)(\tilde{V}) \mid \omega \in N_*\}.$$

Also the second equality of the third point follows straightforwardly: if $x \in B(\mathcal{L}^2(N))$ and

$$\tilde{W}^*(1 \otimes x)\tilde{W} = 1 \otimes x,$$

then $x(\omega \otimes \iota)(\tilde{W}) = (\omega \otimes \iota)(\tilde{W})x$ for all $\omega \in (N^{\text{op}})_*$. From Proposition 1.10.2, we conclude that $x \in \hat{N}'$. As $\tilde{W} \in N^{\text{op}} \otimes \hat{N}$, it is also clear that any $x \in \hat{N}'$ satisfies $\tilde{W}^*(1 \otimes x)\tilde{W} = 1 \otimes x$.

□

Recall that we had introduced in Definition 1.5 the notion of the co-opposite Galois co-object

$$(N^{\text{cop}}, \Delta_{N^{\text{cop}}}) = (N, \Delta_N^{\text{op}}).$$

The following lemma gathers some transfer results between this structure and the original one.

Lemma 1.18. 1. The right regular $(N, \Delta_N^{\text{op}})$ -corepresentation for $(M, \Delta_M^{\text{op}})$ equals $\Sigma \tilde{W}^* \Sigma$, while the left regular $(N, \Delta_N^{\text{op}})$ -corepresentation equals $\Sigma \tilde{V}^* \Sigma$.

2. The dual von Neumann algebra $(N^{\text{cop}})^\wedge$ equals \hat{N}' .

Proof. The two statements are easily verified (the second one follows from Lemma 1.17.3). \square

One can also relate the two adjoint coactions on respectively \widehat{N} and \widehat{N}' , but this result requires some more preparation. We will relegate this investigation to the end of the third section (see Proposition 3.11).

Let us end this section with some remarks on 2-cocycles.

Definition 1.19. ([12]) *Let (M, Δ_M) be a von Neumann bialgebra. A unitary element $\Omega \in M \overline{\otimes} M$ is called a unitary 2-cocycle if Ω satisfies the following identity, called the 2-cocycle identity:*

$$(\Omega \otimes 1)(\Delta_M \otimes \iota)(\Omega) = (1 \otimes \Omega)(\iota \otimes \Delta_M)(\Omega).$$

Example 1.20. *Let (M, Δ_M) be a compact Woronowicz algebra, and Ω a unitary 2-cocycle for (M, Δ_M) . Then if we put $\mathcal{L}^2(N) = \mathcal{L}^2(M)$, $N = M$ and*

$$\Delta_N(x) = \Omega \Delta_M(x), \quad \text{for all } x \in M,$$

the couple (N, Δ_N) is a Galois co-object for (M, Δ_M) , called the Galois co-object associated with Ω .

It is easy to see that if Ω_1 and Ω_2 are two unitary 2-cocycles for (M, Δ_M) , then their associated Galois co-objects are isomorphic iff the unitary 2-cocycles are coboundary equivalent, that is, iff there exists a unitary $u \in M$ such that

$$\Omega_2 = (u^* \otimes u^*) \Omega_1 \Delta_M(u).$$

In particular, the Galois co-object associated with a 2-cocycle Ω on (M, Δ_M) is isomorphic to (M, Δ_M) as a right Galois co-object iff the 2-cocycle is a *coboundary*, i.e. is coboundary equivalent to $1 \otimes 1$.

Definition 1.21. *Let (M, Δ_M) be a compact von Neumann algebra, and (N, Δ_N) a Galois co-object for (M, Δ_M) . Then (N, Δ_N) is called cleft if there exists a unitary 2-cocycle Ω for (M, Δ_M) such that (N, Δ_N) is isomorphic to the Galois co-object associated with Ω .*

At the moment, we do not have any examples of non-cleft Galois co-objects for compact Woronowicz algebras, although these *do* exist in the non-compact case. For example, in [4], non-cleft Galois co-objects were (implicitly) constructed for discrete Woronowicz algebras (see Definition 2.1), the Galois co-object being an l^∞ -direct sum of *rectangular* matrix blocks. For *commutative* compact Woronowicz algebras, that is, those arising from compact groups, it *can* be proven that all Galois co-objects are necessarily cleft (that is, arise from a unitary (measurable) 2-cocycle function on the compact group). We will later prove that this is also the case for *co-commutative* compact Woronowicz algebras (i.e. group von Neumann algebras of discrete groups).

2 Galois objects for discrete Woronowicz algebras

In this section, we will make the connection with the theory of Galois objects from [10].

We first introduce the notion of the dual of a compact Woronowicz algebra.

Definition 2.1. *Let (M, Δ_M) be a compact Woronowicz algebra with regular left corepresentation W . Define*

$$\widehat{M} = \{(\omega \otimes \iota)(W) \mid \omega \in M_*\}^{\sigma\text{-weak closure}}.$$

Then \widehat{M} is a von Neumann algebra which can be endowed with a von Neumann bialgebra structure by giving it the unique comultiplication $\Delta_{\widehat{M}}$ such that

$$(\iota \otimes \Delta_{\widehat{M}})(W) = W_{13}W_{12}.$$

We will call the couple $(\widehat{M}, \Delta_{\widehat{M}})$ the discrete Woronowicz algebra dual to (M, Δ_M) .

In fact, we had already introduced the notation \widehat{M} in the remark after Proposition 1.10, as it can be considered to be the space \widehat{N} in the special case where the right Galois co-object (N, Δ_N) equals (M, Δ_M) . We then also remind that we had introduced some special notations for this case in the Notation 1.14. The following proposition gathers some useful information which can be found in the literature (for example, see the Remark 1.15 in [30], although we warn the reader that their comultiplication on \widehat{M} is opposite to ours).

Proposition 2.2. *Let (M, Δ_M) be a compact Woronowicz algebra, and $(\widehat{M}, \Delta_{\widehat{M}})$ its dual.*

1. *For all $r \in I_M$, the number $m_r = \dim(\mathcal{X}_r)$ is finite.*
2. *There exists a left Δ_M -invariant nsf weight $\varphi_{\widehat{M}}$ on \widehat{M} : for all normal states on \widehat{M} and all positive $x \in \widehat{M}^+$, we have*

$$\varphi_{\widehat{M}}((\omega \otimes \iota)\Delta_{\widehat{M}}(x)) = \varphi_{\widehat{M}}(x).$$

A concrete formula for $\varphi_{\widehat{M}}$ is given by

$$\varphi_{\widehat{M}}(e_{r,ij}) = \delta_{i,j} D_{r,j}^{-1}, \quad \text{for all } r \in I_M, 0 \leq i, j < m_r.$$

3. *On the other hand, define $\psi_{\widehat{M}}$ to be the unique nsf weight on \widehat{M} such that*

$$\psi_{\widehat{M}}(e_{r,ij}) = \delta_{i,j} c_r^2 D_{r,i},$$

where $c_r = \text{Tr}(D_r^{-1})^{1/2}$ (which is known as the quantum dimension of the irreducible corepresentation corresponding to the index $r \in I_M$). Then $\psi_{\widehat{M}}$ is right $\Delta_{\widehat{M}}$ -invariant: for all normal states on \widehat{M} and all positive $x \in \widehat{M}^+$, we have

$$\psi_{\widehat{M}}((\iota \otimes \omega)\Delta_{\widehat{M}}(x)) = \psi_{\widehat{M}}(x).$$

4. *The Radon-Nikodym derivative between $\psi_{\widehat{M}}$ and $\varphi_{\widehat{M}}$ is given by the (possibly unbounded) positive, non-singular operator*

$$\delta_{\widehat{M}} = \bigoplus_{r \in I_M} c_r^2 D_r^2,$$

and $\delta_{\widehat{M}}$ is then a group-like element: for all $t \in \mathbb{R}$, we have

$$\Delta_{\widehat{M}}(\delta_{\widehat{M}}^{it}) = \delta_{\widehat{M}}^{it} \otimes \delta_{\widehat{M}}^{it}.$$

The purpose of this section is to show that for an *arbitrary* Galois co-object (N, Δ_N) , the comultiplication $\Delta_{\widehat{M}}$ can be generalized to a coaction $\alpha_{\widehat{N}}$ of $(\widehat{M}, \Delta_{\widehat{M}})$ on \widehat{N} . This coaction then shares many properties with the actual comultiplication $\Delta_{\widehat{M}}$.

For the rest of this section, we again fix a compact Woronowicz algebra (M, Δ_M) and a right Galois co-object (N, Δ_N) for it. We keep using the notation from the previous section.

Notation 2.3. *We denote by $\varphi_{\widehat{N}}$ the nsf weight on \widehat{N} which is uniquely determined by the fact that all $e_{r,ij} \in \mathcal{M}_{\varphi_{\widehat{N}}}$, with*

$$\varphi_{\widehat{N}}(e_{r,ij}) = \delta_{i,j} T_{r,j}^{-1},$$

where the $T_{r,j}$ were introduced in Notation 1.14. We will then take the GNS-construction for $\varphi_{\widehat{N}}$ also inside $\bigoplus_{r \in I_N} \mathcal{H}_r \otimes \overline{\mathcal{H}_r}$, the GNS-map $\Lambda_{\widehat{N}}$ of $\varphi_{\widehat{N}}$ being determined by

$$\Lambda_{\widehat{N}}(e_{r,ij}) = T_{r,j}^{-1/2} e_{r,i} \otimes \overline{e_{r,j}}.$$

The same notation will be used when (N, Δ_N) equals (M, Δ_M) considered as a right Galois co-object over itself, taking however into consideration the special notations from Notation 1.14.

Remarks:

1. The fact that there exists a *unique* nsf weight with the above properties requires in fact a small technical argument (at least in case the \mathcal{H}_r are not finite-dimensional). The main observations to make are the well-known fact that any nsf weight ψ on a type I -factor is of the form $\text{Tr}(S^{1/2} \cdot S^{1/2})$ for some non-singular positive (possibly unbounded) operator S (see [27], Lemma VIII.2.8), and the fact that if ξ is a vector with $\psi(\theta_{\xi,\xi}) < \infty$ (where we recall that $\theta_{\xi,\xi}$ is the rank one operator associated with ξ), then $\xi \in \mathcal{D}(S^{1/2})$ with $\|S^{1/2}\xi\|^2 = \psi(\theta_{\xi,\xi})$ (this can, for example, be pieced together from the results in [27], section IX.3). With this information, it should then be easy to verify that the nsf weight $\varphi_{\hat{N}}$ in the previous notation is indeed well-defined and uniquely determined.
2. It is easy to check, using the orthogonality relations between the $\widetilde{W}_{r,ij}$, that for $r \in I_N$ and $0 \leq i, j < n_r$, we have

$$(\varphi_M(\cdot \widetilde{W}_{r,ij}^*) \otimes \iota)(\widetilde{W}) \in \mathcal{N}_{\varphi_{\hat{N}}},$$

with

$$\Lambda_{\hat{N}}((\varphi_M(\cdot \widetilde{W}_{r,ij}^*) \otimes \iota)(\widetilde{W})) = \widetilde{W}_{r,ij}^* \xi_M.$$

Hence our identifications of $\mathcal{L}^2(N)$ and $\mathcal{L}^2(\hat{N})$ coincide with the ‘usual’ way in which Pontryagin duality is defined in the setting of (locally) compact quantum groups (see [20]).

Proposition 2.4. *Denote by $\alpha_{\hat{N}}$ the unital normal faithful $*$ -homomorphism*

$$\alpha_{\hat{N}} : \hat{N} \rightarrow \hat{N} \bar{\otimes} B(\mathcal{L}^2(M)) : x \rightarrow \Sigma \widetilde{W}(x \otimes 1) \widetilde{W}^* \Sigma.$$

Then $\alpha_{\hat{N}}$ has range in $\hat{N} \bar{\otimes} \widehat{M}$, and determines an ergodic coaction of $(\widehat{M}, \Delta_{\widehat{M}})$ on \hat{N} .

Before giving the proof, we first state a lemma that we will need in the course of it.

Lemma 2.5. *Take $r \in I_N$ arbitrary. Then the linear span of the set*

$$\{\widetilde{W}_{r,ij}^* a \mid 0 \leq i, j < n_r, a \in M\}$$

is σ -weakly dense in N .

Proof. Denote by \widetilde{N} the σ -weak closure of the linear span of $\{\widetilde{W}_{r,ij}^* a \mid 0 \leq i, j < n_r, a \in M\}$. Then clearly $\widetilde{N} \subseteq N$. Now choose $r \in I_N$ fixed, and take an $x \in N$. By Proposition 1.15.2, we have

$$x = \sum_{k=0}^{n_r-1} \widetilde{W}_{r,k0}^* (\widetilde{W}_{r,k0} x),$$

the sum converging σ -weakly. As $\widetilde{W}_{r,k0} x \in M$ for all $0 \leq k < n_r$, the sum on the right hand side lies in \widetilde{N} . So also $N \subseteq \widetilde{N}$. □

Proof (of Proposition 2.4). Take $\omega \in (N^{\text{op}})_*$, and denote $x = (\omega \otimes \iota)(\widetilde{W})$. By the pentagonal identity for \widetilde{W} (Proposition 1.7.4), we easily get that

$$\alpha_{\hat{N}}(x) = (\omega \otimes \iota \otimes \iota)(W_{13} \widetilde{W}_{12}) \in \hat{N} \bar{\otimes} \widehat{M},$$

and by an application of the formula $(\iota \otimes \Delta_{\widehat{M}})(W) = W_{13} W_{12}$, we find

$$(\alpha_{\hat{N}} \otimes \iota) \alpha_{\hat{N}}(x) = (\iota \otimes \Delta_{\widehat{M}}) \alpha_{\hat{N}}(x).$$

As elements of the form x constitute a σ -weakly dense subspace of \widehat{N} by Proposition 1.10.2, we have proven that $\alpha_{\widehat{N}}$ is a well-defined coaction.

We now show that it is ergodic. Take an element $x \in \widehat{N}$ satisfying $\alpha_{\widehat{N}}(x) = x \otimes 1$. Then for all $y \in \widehat{N}$, we get

$$(x \otimes 1)\text{Ad}_R(y) = \text{Ad}_R(y)(x \otimes 1),$$

where we recall that $\text{Ad}_R(y) = \Sigma \widetilde{W}(1 \otimes y) \widetilde{W}^* \Sigma$. As $\{(\iota \otimes \omega)\text{Ad}_R(y) \mid \omega \in M_*\}'' = \widehat{N}$, a general fact for any coaction of a compact Woronowicz algebra, we find $x \in \mathcal{Z}(\widehat{N})$, the center of \widehat{N} .

Write then $x = \sum_{r \in I_N} x_r p_r$, where $r \rightarrow x_r \in l^\infty(I_N)$ and p_r the r -th minimal central projection of \widehat{N} . Then as $\alpha_{\widehat{N}}(x) = x \otimes 1$, we have

$$x(\iota \otimes \omega)(\widetilde{W}^* \Sigma) = (\iota \otimes \omega)(\widetilde{W}^* \Sigma)x$$

for all $\omega \in B(\mathcal{L}^2(M), \mathcal{L}^2(N))_*$. If we take $\omega = \omega_{a\xi_M, \widetilde{W}_{r,ij}^* \xi_M}$ for some $r \in I_N$, $0 \leq i, j < n_r$ and $a \in M$, and apply both sides of the above equality to $\widetilde{W}_{r,kj}^* \xi_M$ for some $0 \leq k < n_r$, we get, by using the orthogonality relations for the $\widetilde{W}_{r,ij}$, that

$$x \widetilde{W}_{r,ki}^* a \xi_M = x_r \widetilde{W}_{r,ki}^* a \xi_M.$$

As, with r a *fixed* element of I_N , we have that $\{\widetilde{W}_{r,ki}^* a \mid a \in M\}$ is σ -weakly dense in N , by the previous lemma, we get that $x\xi = x_r \xi$ for all $\xi \in \mathcal{L}^2(N)$, so x is a scalar multiple of the unit. \square

Our next goal is to show that $\alpha_{\widehat{N}}$ is nicely behaved with respect to the weight structure on $(\widehat{M}, \Delta_{\widehat{M}})$.

Proposition 2.6. *For all $x \in \widehat{N}^+$ and all normal positive states ω on \widehat{N} , we have*

$$\varphi_{\widehat{M}}((\omega \otimes \iota)\alpha_{\widehat{N}}(x)) = \varphi_{\widehat{N}}(x).$$

In particular, $\alpha_{\widehat{N}}$ is an integrable coaction.

Remark: The fact that $\alpha_{\widehat{N}}$ is integrable means that there exists a σ -weakly dense subspace of \widehat{N} consisting of elements x for which *all* expressions $(\omega \otimes \iota)\alpha_{\widehat{N}}(x)$ with $\omega \in \widehat{N}_*$ lie in $\mathcal{M}_{\varphi_{\widehat{M}}}$.

Proof. We first recall a small technical result from [29], Proposition 1.3. Namely, as $\alpha_{\widehat{N}}$ is an ergodic coaction, there exists a (not necessarily semi-finite) normal faithful weight $\varphi'_{\widehat{N}}$ on \widehat{N} , determined by the following formula: for all $x \in \widehat{N}_+$, we have

$$\varphi'_{\widehat{N}}(x) = \varphi_{\widehat{M}}((\omega \otimes \iota)\alpha_{\widehat{N}}(x)),$$

where ω is *any* normal state on \widehat{N} . Our job then is to prove that $\varphi'_{\widehat{N}} = \varphi_{\widehat{N}}$. By the remark after Notation 2.3, it is enough to prove that the $e_{r,ij}$ are in $\mathcal{M}_{\varphi'_{\widehat{N}}}$ with

$$\varphi'_{\widehat{N}}(e_{r,ij}) = \delta_{i,j} T_{r,j}^{-1}.$$

Take $r, s \in I_N$ and $0 \leq i, j < n_r$, $0 \leq k, l < n_s$. Then we compute, using the GNS-construction for $\varphi_{\widehat{N}}$ from Notation 2.3 and the functionals $\omega_{s,kl}$ introduced in Notation 1.14, that

$$\begin{aligned}
& (\iota \otimes \omega_{s,kl})(\widetilde{W})\Lambda_{\widehat{N}}(e_{r,ij}) \\
&= T_{r,j}^{-1} \widetilde{W}_{s,lk} \widetilde{W}_{r,ij}^* \xi_M \\
&= T_{r,j}^{-1} \sum_{t \in I_M} \sum_{m,n=0}^{m_t-1} \langle \widetilde{W}_{s,lk} \widetilde{W}_{r,ij}^* \xi_M, \frac{1}{\|\widetilde{W}_{t,mn}^* \xi_M\|} W_{t,mn}^* \xi_M \rangle \frac{1}{\|\widetilde{W}_{t,mn}^* \xi_M\|} W_{t,mn}^* \xi_M \\
&= T_{r,j}^{-1} \sum_{t \in I_M} \sum_{m,n=0}^{m_t-1} \varphi_M(W_{t,mn} \widetilde{W}_{s,lk} \widetilde{W}_{r,ij}^*) D_{t,n}^{-1} W_{t,mn}^* \xi_M \\
&= T_{r,j}^{-1} \sum_{t \in I_M} \sum_{m,n=0}^{m_t-1} \varphi_M(W_{t,mn} \widetilde{W}_{s,lk} \widetilde{W}_{r,ij}^*) \Lambda_{\widehat{M}}(e_{t,mn}),
\end{aligned}$$

the latter sums converging in norm.

On the other hand,

$$\begin{aligned}
(\omega_{s,kl} \otimes \iota)\alpha_{\widehat{N}}(e_{r,ij}) &= T_{r,j}^{-1}(\omega_{s,kl} \otimes \iota)\alpha_{\widehat{N}}((\varphi_M(\cdot \widetilde{W}_{r,ij}^*) \otimes \iota)(\widetilde{W})) \\
&= T_{r,j}^{-1}(\varphi_M(\cdot \widetilde{W}_{r,ij}^*) \otimes \omega_{s,kl} \otimes \iota)(W_{13} \widetilde{W}_{12}) \\
&= T_{r,j}^{-1} \sum_{t \in I_M} \sum_{m,n=0}^{m_t-1} \varphi_M(W_{t,mn} \widetilde{W}_{s,lk} \widetilde{W}_{r,ij}^*) e_{t,mn},
\end{aligned}$$

where the latter sum now converges in the strong topology.

As $\Lambda_{\widehat{M}}$ is a strong-norm closed map from $\mathcal{N}_{\varphi_{\widehat{M}}}$ to $\mathcal{L}^2(\widehat{M})$, it follows that

$$(\omega_{s,kl} \otimes \iota)(\alpha_{\widehat{N}}(e_{r,ij})) \in \mathcal{N}_{\varphi_{\widehat{M}}},$$

with

$$\Lambda_{\widehat{M}}((\omega_{s,kl} \otimes \iota)(\alpha_{\widehat{N}}(e_{r,ij}))) = (\iota \otimes \omega_{s,kl})(\widetilde{W})\Lambda_{\widehat{N}}(e_{r,ij}).$$

Again by closedness, these assertions remain true when $\omega_{s,kl}$ is replaced by an arbitrary normal functional on \widehat{N} .

Let now $\xi_{r,i,j} = e_{r,i} \otimes \overline{e_{r,j}}$ for $r \in I_N$ and $0 \leq i, j < n_r$. For any finite subset \mathcal{J}_0 of the set $\mathcal{J} = \{(r, i, j) \mid r \in I_N, 0 \leq i, j < n_r\}$, denote by $P_{\mathcal{J}_0}$ the orthogonal projection onto the linear span of the ξ_n with $n \in \mathcal{J}_0$. Take an arbitrary state $\omega \in \widehat{N}_*$ and $x = y^*y$ in the linear span of the $e_{r,ij}$. We remark then that there exists a unit vector $\xi \in \mathcal{L}^2(\widehat{N})$ with $\omega = \omega_{\xi,\xi} (= \langle \cdot, \xi \rangle)$. We can now compute, using the normality of our weights, that

$$\begin{aligned}
\varphi_{\widehat{M}}((\omega \otimes \iota)\alpha_{\widehat{N}}(x)) &= \lim_{\substack{\mathcal{J}_0 \subseteq \mathcal{J} \\ \text{fin}}} \varphi_{\widehat{M}}((\omega_{\xi,\xi} \otimes \iota)(\alpha_{\widehat{N}}(y)^*(P_{\mathcal{J}_0} \otimes 1)\alpha_{\widehat{N}}(y))) \\
&= \lim_{\substack{\mathcal{J}_0 \subseteq \mathcal{J} \\ \text{fin}}} \sum_{n \in \mathcal{J}_0} \varphi_{\widehat{M}}((\omega_{\xi,\xi_n} \otimes \iota)(\alpha_{\widehat{N}}(y))^* \cdot (\omega_{\xi,\xi_n} \otimes \iota)(\alpha_{\widehat{N}}(y))) \\
&= \lim_{\substack{\mathcal{J}_0 \subseteq \mathcal{J} \\ \text{fin}}} \sum_{n \in \mathcal{J}_0} \|(\iota \otimes \omega_{\xi,\xi_n})(\widetilde{W})\Lambda_{\widehat{N}}(y)\|^2 \\
&= \varphi_{\widehat{N}}(y^*y) = \varphi_{\widehat{N}}(x),
\end{aligned}$$

by the unitarity of \widetilde{W} . From this, it immediately follows that all $e_{r,ij}$ are integrable for $\varphi'_{\widehat{N}}$, and that $\varphi'_{\widehat{N}} = \varphi_{\widehat{N}}$ on the linear span of the $e_{r,ij}$. This then concludes the proof. \square

We have shown so far that $\alpha_{\widehat{N}}$ is an integrable, ergodic coaction. The final property of $\alpha_{\widehat{N}}$ is that a certain isometry which can be constructed from $\alpha_{\widehat{N}}$ is in fact a unitary.

Proposition 2.7. *Take $x, y \in \mathcal{N}_{\varphi_{\widehat{N}}}$. Then $\alpha_{\widehat{N}}(y)(x \otimes 1) \in \mathcal{N}_{\varphi_{\widehat{N}} \otimes \varphi_{\widehat{M}}}$, and*

$$(\Lambda_{\widehat{N}} \otimes \Lambda_{\widehat{M}})(\alpha(y)(x \otimes 1)) = \Sigma \widetilde{W} \Sigma \Lambda_{\widehat{N}}(x) \otimes \Lambda_{\widehat{N}}(y).$$

Proof. The claim concerning the square integrability of $\alpha_{\widehat{N}}(y)(x \otimes 1)$ follows immediately from the fact that $\alpha_{\widehat{N}}$ is integrable, with $(\iota \otimes \varphi_{\widehat{M}})\alpha_{\widehat{N}} = \varphi_{\widehat{N}}$. Moreover, we can then define an isometry

$$\widetilde{G} : \mathcal{L}^2(N) \otimes \mathcal{L}^2(N) \rightarrow \mathcal{L}^2(M) \otimes \mathcal{L}^2(N)$$

inside $B(\mathcal{L}^2(N), \mathcal{L}^2(M)) \widehat{\otimes} \widehat{N}$ such that precisely

$$(\Lambda_{\widehat{N}} \otimes \Lambda_{\widehat{M}})(\alpha(y)(x \otimes 1)) = \Sigma \widetilde{G} \Sigma \Lambda_{\widehat{N}}(x) \otimes \Lambda_{\widehat{N}}(y)$$

for all $x, y \in \mathcal{N}_{\varphi_{\widehat{N}}}$. We need to show that $\widetilde{G} = \widetilde{W}$.

However, it is easily seen that for all $\omega \in B(\mathcal{L}^2(\widehat{N}))_*$ and $x \in \mathcal{N}_{\varphi_{\widehat{N}}}$, we will have

$$(\iota \otimes \omega)(\widetilde{G})\Lambda_{\widehat{N}}(x) = \Lambda_{\widehat{M}}((\omega \otimes \iota)\alpha_{\widehat{N}}(x)).$$

By the computations made in the previous proposition, it follows that $(\iota \otimes \omega)(\widetilde{G})$ coincides with $(\iota \otimes \omega)(\widetilde{W})$ on the linear span of the $e_{r,i} \otimes \overline{e}_{r,j}$ for $r \in I_N$, $0 \leq i, j < n_r$, and hence $\widetilde{G} = \widetilde{W}$. \square

The three propositions above immediately show the following.

Theorem 2.8. *Let (M, Δ_M) be a compact Woronowicz algebra, (N, Δ_N) a right Galois co-object for (M, Δ_M) . Then the couple $(\widehat{N}, \alpha_{\widehat{N}})$ makes \widehat{N} into a right Galois object for the discrete Woronowicz algebra $(\widehat{M}, \Delta_{\widehat{M}})$, with corresponding Galois unitary $\widetilde{G} = \widetilde{W}$.*

For the terminology ‘Galois object’, we refer the reader to [8] (where the notations N and \widehat{N} are interchanged). In fact, it is simply *defined* to be an integrable ergodic coaction for which the map \widetilde{G} , as we constructed it in the course of the proof the previous proposition, is a unitary. This map \widetilde{G} is in general called the *Galois unitary* associated with the Galois object, and as we saw in the previous proposition, it coincides precisely with \widetilde{W} in case the Galois object is constructed from a Galois co-object for a compact Woronowicz algebra. Galois objects can also be defined as being *ergodic, semi-dual coactions* (see [29], Proposition 5.12 for the terminology, and the remark under Proposition 3.5 of [8] for the connection). We further remark that Galois objects for *compact* Woronowicz algebras were treated in [4] (where they are termed ‘actions of full quantum multiplicity’), and for *commutative* compact Woronowicz algebras, that is for ordinary compact groups, in [33] and [21] (where they are termed ‘actions of full multiplicity’). For Galois objects in the Hopf algebra setting, we refer to the overview [26].

We may now use the results from [8], which we gather in the following theorem.

Theorem 2.9. *Let (N, Δ_N) be a right Galois co-object for a compact Woronowicz algebra (M, Δ_M) , and let $(\widehat{N}, \alpha_{\widehat{N}})$ be the associated right Galois object for the dual discrete Woronowicz algebra $(\widehat{M}, \Delta_{\widehat{M}})$. Then the following statements hold.*

1. *There exists an nsf weight $\psi_{\widehat{N}}$ on \widehat{N} , unique up to scaling with a positive constant, which is $\alpha_{\widehat{N}}$ -invariant: for all states $\omega \in \widehat{M}_*$ and all $x \in \widehat{N}^+$, we have*

$$\psi_{\widehat{N}}((\iota \otimes \omega)\alpha_{\widehat{N}}(x)) = \psi_{\widehat{N}}(x).$$

2. The Radon-Nikodym derivative $\delta_{\widehat{N}}\eta_{\widehat{N}}$ of $\psi_{\widehat{N}}$ with respect to $\varphi_{\widehat{N}}$ satisfies

$$\alpha_{\widehat{N}}(\delta_{\widehat{N}}^{it}) = \delta_{\widehat{N}}^{it} \otimes \delta_M^{it} \quad \text{for all } t \in \mathbb{R}.$$

3. The Radon-Nikodym derivative $\delta_{\widehat{N}}$ is $\sigma_t^{\varphi_{\widehat{N}}}$ -invariant, where the latter denotes the modular one-parameter group associated with $\varphi_{\widehat{N}}$:

$$\sigma_t^{\varphi_{\widehat{N}}}(\delta_{\widehat{N}}^{is}) = \delta_{\widehat{N}}^{is} \quad \text{for all } s, t \in \mathbb{R}.$$

Proof. See [8], Theorem 3.18, Proposition 3.15 and Lemma 3.17. \square

Notation 2.10. We denote by A_r the non-singular (possibly unbounded) positive operator $p_r \delta_{\widehat{N}} \in B(\mathcal{H}_r)$, so that

$$\delta_{\widehat{N}} = \bigoplus_{r \in I_N} A_r.$$

By the third item in the previous theorem, the operator A_r strongly commutes with T_r . In particular, this means that A_r is diagonalizable, and that we may choose our $e_{r,i} \in \mathcal{H}_r$ so that they are *also* eigenvectors for the A_r . We then write $A_{r,i}$ for the eigenvalue of A_r with respect to the eigenvector $e_{r,i}$.

Remark: Let \mathcal{A} be the Hopf $*$ -algebra associated with (M, Δ_M) , consisting of all elements $x \in M$ with $\Delta_M(x) \in M \odot M$, the algebraic tensor product. Then it is well-known that \mathcal{A} is a σ -weakly dense sub- $*$ -algebra of M , closed under the modular automorphism group $\sigma_t^{\varphi_M}$ of φ_M . Let \mathcal{B}_r be the sub- $*$ -algebra of $B(\mathcal{H}_r)$ consisting of all elements $x \in B(\mathcal{H}_r)$ with $\text{Ad}_R^{(r)}(x) \in B(\mathcal{H}_r) \odot \mathcal{A}$. Again, it is well-known that \mathcal{B}_r is a σ -weakly dense sub- $*$ -algebra of $B(\mathcal{H}_r)$ (it is the linear span of the coefficients of the spectral subspaces associated with $\text{Ad}_R^{(r)}$, see for example [4]). Then the operators A_r , introduced in the above notation, turn out to be determined, up to a scalar, by the formula

$$A_r^{it} x A_r^{-it} = (\iota \otimes \varepsilon \circ \sigma_t^{\varphi_M}) \text{Ad}_R^{(r)}(x), \quad \text{for all } x \in \mathcal{B}_r,$$

where ε denotes the counit of \mathcal{A} . This formula can be derived from the way in which $\delta_{\widehat{N}}$ was constructed in [8]. Hence, up to multiplication with a non-singular (possibly unbounded) positive element in the center of \widehat{N} , the operator $\delta_{\widehat{N}}$ can be recovered from the knowledge of all the $\text{Ad}_R^{(r)}$.

3 Projective representations of compact quantum groups

Using the results from the first section, we can easily develop a Peter-Weyl theory for projective representations of compact quantum groups. We will in the following use again the notation which we introduced in the first section.

We first define the notion of a projective representation relative to a fixed Galois co-object.

Definition 3.1. Let (M, Δ_M) be a compact Woronowicz algebra, (N, Δ_N) a right Galois co-object for (M, Δ_M) . A (left) (N, Δ_N) -corepresentation of (M, Δ_M) on a Hilbert space \mathcal{H} consists of a unitary map $\mathcal{G} \in N \overline{\otimes} B(\mathcal{H})$ such that

$$(\Delta_N \otimes \iota)\mathcal{G} = \mathcal{G}_{13}\mathcal{G}_{23}.$$

We call a Hilbert subspace $\mathcal{K} \subseteq \mathcal{H}$ invariant w.r.t. the (N, Δ_N) -corepresentation when \mathcal{G} restricts to a unitary in $N \overline{\otimes} B(\mathcal{K})$.

We call \mathcal{G} irreducible if the only invariant Hilbert subspaces are 0 and \mathcal{H} , and indecomposable when \mathcal{H} can not be written as the direct sum of two non-zero invariant subspaces.

We call two (N, Δ_N) -corepresentations \mathcal{G}_1 and \mathcal{G}_2 on respective Hilbert spaces \mathcal{H}_1 and \mathcal{H}_2 unitarily equivalent if there exists $u \in B(\mathcal{H}_1, \mathcal{H}_2)$ such that

$$\mathcal{G}_2(1 \otimes u) = (1 \otimes u)\mathcal{G}_1.$$

Remark: When (N, Δ_N) comes from a 2-cocycle Ω for (M, Δ_M) , we will also simply speak of Ω -corepresentations.

Theorem 3.2. *Let (M, Δ_M) be a compact quantum group, (N, Δ_N) a right Galois co-object for (M, Δ_M) . Denote by \tilde{V} the right regular (N, Δ_N) -corepresentation for (M, Δ_M) , and let*

$$\tilde{V}_r = (p_r \otimes 1)\tilde{V} \in N \bar{\otimes} B(\overline{\mathcal{H}_r})$$

be the components of \tilde{V} , where the p_r denote the minimal projections of $\mathcal{Z}(\hat{N})$.

1. The unitaries $\Sigma \tilde{V}_r \Sigma$ are indecomposable left (N, Δ_N) -corepresentations on the Hilbert spaces $\overline{\mathcal{H}_r}$.
2. Any indecomposable (N, Δ_N) -corepresentation is unitarily equivalent with a unique $\Sigma \tilde{V}_r \Sigma$.
3. Any (N, Δ_N) -corepresentation splits as a direct sum of indecomposable (N, Δ_N) -corepresentations.
4. Any indecomposable (N, Δ_N) -corepresentation is irreducible.

Proof. As $(\iota \otimes \Delta_N)\tilde{V} = \tilde{V}_{12}\tilde{V}_{13}$, we immediately get that the unitaries $\Sigma \tilde{V}_r \Sigma$ are left (N, Δ_N) -corepresentations. By Lemma 1.17.3, the space

$$\{(\iota \otimes \omega)(\tilde{V}_r) \mid \omega \in B(\mathcal{L}^2(M), \mathcal{L}^2(N))_*\}$$

equals the whole of $B(\overline{\mathcal{H}_r})$, from which it immediately follows that $\Sigma \tilde{V}_r \Sigma$ is indecomposable, and even irreducible.

For the second statement, we use that the linear span of the matrix entries of the \tilde{V}_r 's span a norm-dense subset of $\mathcal{L}^2(N)$ when applied to ξ_M . Hence, if \mathcal{G} is an indecomposable left (N, Δ_N) -corepresentation of (M, Δ_M) on a Hilbert space \mathcal{H} , there must exist some $r \in I_N$ and an $x \in B(\mathcal{H}, \overline{\mathcal{H}_r})$ such that

$$(\varphi_M \otimes \iota)((\Sigma \tilde{V}_r \Sigma)^*(1 \otimes x)\mathcal{G}) \neq 0.$$

As in the proof of Theorem 1.15, this forces a scalar multiple of this expression to be a unitary intertwiner, proving that \mathcal{G} is isomorphic to $\Sigma \tilde{V}_r \Sigma$. By the orthogonality relations between the $\Sigma \tilde{V}_r \Sigma$, these are all pairwise non-isomorphic. Hence the above r for \mathcal{G} is uniquely determined.

To prove the third statement, consider the normal faithful unital *-homomorphism

$$\alpha : B(\mathcal{H}) \rightarrow M \bar{\otimes} B(\mathcal{H}) : x \rightarrow \mathcal{G}^*(1 \otimes x)\mathcal{G}.$$

As $(\Delta_N \otimes \iota)\mathcal{G} = \mathcal{G}_{13}\mathcal{G}_{23}$, we see that α is a coaction. Let $Z = B(\mathcal{H})^\alpha$, the set of fixed points for α . As in the proof of Proposition 1.10, we have that Z is the range of a normal conditional expectation on $B(\mathcal{H})$. Hence Z is a von Neumann algebraic direct sum of type I -factors. Let then A be an atomic maximal abelian von Neumann subalgebra of Z , and denote by p_i the set of minimal projections in A . Then it is clear that each $p_i \mathcal{H}$ is a fixed subspace for \mathcal{G} . The spaces $p_i \mathcal{H}$ must further be indecomposable: for if not, then we could find a p_i and a non-zero projection p in $B(\mathcal{H})$ with p strictly smaller than p_i and both $p \mathcal{H}$ and $(p_i - p) \mathcal{H}$ invariant under \mathcal{G} . This would imply that p is a fixed

element for α , commuting with all $x \in A$. Hence $p \in A$ by maximal abelianness. As p_i was a minimal projection in A , this gives a contradiction.

As for the fourth point, we may take our indecomposable (N, Δ_N) -corepresentation to be some $\Sigma \tilde{V}_r \Sigma$, for which we have already proven irreducibility in the proof of the first point. □

Corollary 3.3. *Let (N, Δ_N) be a right Galois co-object for a compact Woronowicz algebra (M, Δ_M) . Let \mathcal{G}_i , $i \in I$, be a maximal set of non-isomorphic irreducible (N, Δ_N) -corepresentations on Hilbert spaces \mathcal{H}_i . Then $I \cong I_N$, and $\tilde{N} \cong \bigoplus_{i \in I} B(\mathcal{H}_i)$.*

Proof. This follows immediately from the second point of the previous proposition. □

We can now pass to projective representations without reference to a fixed Galois co-object.

Definition 3.4. *Let (M, Δ_M) be a compact Woronowicz algebra. A projective (left) corepresentation of (M, Δ_M) on a Hilbert space \mathcal{H} consists of a left coaction α of (M, Δ_M) on $B(\mathcal{H})$,*

$$\alpha : B(\mathcal{H}) \rightarrow M \bar{\otimes} B(\mathcal{H}).$$

Remarks:

1. Interpreting (M, Δ_M) as the space of \mathcal{L}^∞ -functions on some ‘compact quantum group’ \mathbb{G} , the above then corresponds to having a (necessarily continuous) action of \mathbb{G} on $B(\mathcal{H})$. As $\text{Aut}(B(\mathcal{H})) \cong \mathcal{U}(\mathcal{H})/S^1$, this indeed captures the notion of a projective representation when \mathbb{G} is an actual compact group.
2. One similarly has the notion of a *projective right corepresentation* of (M, Δ_M) , for which we replace the left coaction α above by a right coaction. For example, the coactions Ad_R and $\text{Ad}_R^{(r)}$ from the first section are then projective right corepresentations. At the end of this section, we will show one can pass from left to right projective representations, so that one may essentially restrict oneself to the study of projective left corepresentations, as we will do.
3. Some results on (special) coactions of compact Kac algebras on type I factors appear in [22].

In [10], we proved that from any projective corepresentation α , one can construct a Galois co-object (N, Δ_N) together with an (N, Δ_N) -corepresentation \mathcal{G} ‘implementing’ α . We will state the proposition and give a sketch of the proof. For the full proof, we refer the reader to [10].

Proposition 3.5. *Let (M, Δ_M) be a compact Woronowicz algebra, \mathcal{H} a Hilbert space, and α a projective corepresentation of (M, Δ_M) on \mathcal{H} . Then there exists a right Galois co-object (N, Δ_N) for (M, Δ_M) , together with an (N, Δ_N) -corepresentation \mathcal{G} of (M, Δ_M) on \mathcal{H} , such that*

$$\mathcal{G}^*(1 \otimes x)\mathcal{G} = \alpha(x), \quad \text{for all } x \in B(\mathcal{H}).$$

Sketch of proof. Choose a basis $\{e_i\}_{i \in I}$ of \mathcal{H} , and fix an element $0 \in I$. We can then consider the Hilbert space $\mathcal{L}^2(N) = \alpha(e_{00})(\mathcal{L}^2(M) \otimes \mathcal{H})$. We can construct a unitary

$$\mathcal{G} : \mathcal{L}^2(M) \otimes \mathcal{H} \rightarrow \mathcal{L}^2(N) \otimes \mathcal{H} : \xi \rightarrow \sum_{i \in I} (\alpha(e_{0i})\xi) \otimes e_i.$$

Denote by \mathcal{G}_{ij} the i, j -th component of \mathcal{G} . Then \mathcal{G}_{ij} is an operator from $\mathcal{L}^2(M)$ to $\mathcal{L}^2(N)$. We define

$$N = \{\mathcal{G}_{ij}m \mid i, j \in I, m \in M\}^{\sigma\text{-weakly closed linear span}}.$$

It is then possible to construct a map $\Delta_N : N \rightarrow N \bar{\otimes} N$, uniquely determined by the properties that

$$(\Delta_N \otimes \iota)\mathcal{G} = \mathcal{G}_{[13]}\mathcal{G}_{[23]}$$

(where we have added brackets in the leg numbering notation to distinguish them from the indices for matrix coefficients of \mathcal{G}) and

$$\Delta_N(xy) = \Delta_N(x)\Delta_M(y), \quad \text{for all } x \in N, y \in M.$$

One proves that (N, Δ_N) is a Galois co-object for (M, Δ_M) , and then it immediately follows from the above property that \mathcal{G} is a left (N, Δ_N) -corepresentation of (M, Δ_M) on \mathcal{H} . Finally, one proves that $\mathcal{G}^*(1 \otimes x)\mathcal{G} = \alpha(x)$ by direct computation. \square

Definition 3.6. *Let α be a projective corepresentation of a compact Woronowicz algebra (M, Δ_M) on a Hilbert space \mathcal{H} . Denote by (N, Δ_N) the Galois co-object constructed from α as in the above proposition, and denote by $[(N, \Delta_N)]$ its isomorphism class. Then we say that α is an $[(N, \Delta_N)]$ -corepresentation.*

It can be proven (see Proposition 3.4 in [10]) that if α is an $[(N, \Delta_N)]$ -corepresentation of (M, Δ_M) on a Hilbert space \mathcal{H} , and if there exists a Galois co-object $(\tilde{N}, \Delta_{\tilde{N}})$ for (M, Δ_M) which possesses an $(\tilde{N}, \Delta_{\tilde{N}})$ -corepresentation on \mathcal{H} implementing α , then necessarily $[(\tilde{N}, \Delta_{\tilde{N}})] = [(N, \Delta_N)]$. One may regard the isomorphism class of such a Galois co-object as a generalization of the notion of a 2-cohomology class. Also remark that if $\mathcal{G} \in N \bar{\otimes} B(\mathcal{H})$ is a projective (N, Δ_N) -corepresentation of (M, Δ_M) on a Hilbert space \mathcal{H} , then the associated projective corepresentation

$$\alpha : B(\mathcal{H}) \rightarrow M \bar{\otimes} B(\mathcal{H}) : x \rightarrow \mathcal{G}^*(1 \otimes x)\mathcal{G}$$

is an $[(N, \Delta_N)]$ -corepresentation by the above uniqueness result.

If (N, Δ_N) is a Galois co-object for a compact Woronowicz algebra (M, Δ_M) , and an $[(N, \Delta_N)]$ -corepresentation α for (M, Δ_M) is given, then it is in general not true that all (N, Δ_N) -corepresentations implementing α are isomorphic: consider for example ordinary one-dimensional representations. We will see a further instance of this in the final section.

Definition 3.7. *Let (M, Δ_M) be a compact Woronowicz algebra, α a projective corepresentation of (M, Δ_M) on a Hilbert space \mathcal{H} . We say that α is an irreducible projective corepresentation if α is ergodic.*

Proposition 3.8. *Let (N, Δ_N) be a right Galois co-object for a compact Woronowicz algebra (M, Δ_M) , let α be an $[(N, \Delta_N)]$ -corepresentation of (M, Δ_M) on a Hilbert space \mathcal{H} , and let \mathcal{G} be an (N, Δ_N) -corepresentation implementing α .*

Then there is a one-to-one correspondence between the set of α -fixed self-adjoint projections in $B(\mathcal{H})$ and the \mathcal{G} -invariant subspaces \mathcal{K} of \mathcal{H} , given by the correspondence

$$p \rightarrow \mathcal{K} = p\mathcal{H}.$$

In particular, α is irreducible iff \mathcal{G} is irreducible.

Proof. By assumption, we have that

$$\alpha(x) = \mathcal{G}^*(1 \otimes x)\mathcal{G} \quad \text{for all } x \in B(\mathcal{H}).$$

So if p is a self-adjoint projection in $B(\mathcal{H})$ with $\alpha(p) = 1 \otimes p$, we have

$$\mathcal{G}(1 \otimes p) = (1 \otimes p)\mathcal{G},$$

and hence

$$\mathcal{G}(\mathcal{L}^2(M) \otimes p\mathcal{H}) \subseteq \mathcal{L}^2(N) \otimes p\mathcal{H},$$

which means $p\mathcal{H}$ is a \mathcal{G} -invariant subspace.

Conversely, if \mathcal{K} is a \mathcal{G} -invariant subspace, and p the projection onto \mathcal{K} , then also \mathcal{K}^\perp is \mathcal{G} -invariant by Theorem 3.2. Hence we have

$$\mathcal{G}(1 \otimes p) = (1 \otimes p)\mathcal{G},$$

and so $\alpha(p) = 1 \otimes p$, i.e. p is an α -fixed projection. \square

Remark: We in fact already used the above argument in the course of proving Theorem 3.2.3.

Our next proposition shows how projective corepresentations and ordinary corepresentations mesh together.

Proposition 3.9. *Let (N, Δ_N) be a right Galois co-object for a compact Woronowicz algebra (M, Δ_M) . Let \mathcal{G} be a projective left (N, Δ_N) -corepresentation on a Hilbert space \mathcal{H} , and let U be an ordinary left corepresentation of (M, Δ_M) on a Hilbert space \mathcal{K} .*

Then the following statements hold.

1. *The unitary $\mathcal{G}_{12}U_{13} \in N \bar{\otimes} B(\mathcal{H} \otimes \mathcal{K})$ is a unitary (N, Δ_N) -corepresentation on $\mathcal{H} \otimes \mathcal{K}$.*
2. *If both \mathcal{G} and U are irreducible, then $\mathcal{G}_{12}U_{13}$ is a finite direct sum of irreducible (N, Δ_N) -corepresentations.*

Proof. The fact that $\mathcal{G}_{12}U_{13}$ is a unitary (N, Δ_N) -corepresentation is trivial to verify. If further \mathcal{G} and U are irreducible, then we know already that $\mathcal{G}_{12}U_{13} = \bigoplus_{i \in J} \mathcal{G}_i$ for a certain set \mathcal{G}_i of irreducible (N, Δ_N) -corepresentations indexed by a parameter set J . We have to prove that J is finite.

Let α be the projective corepresentation associated with \mathcal{G} , so

$$\alpha(x) = \mathcal{G}^*(1 \otimes x)\mathcal{G}, \quad x \in B(\mathcal{H}).$$

By the previous proposition, we know that α is ergodic. Let \mathcal{B} be the linear span of the spectral subspaces inside $B(\mathcal{H})$, which is a σ -weakly dense sub- $*$ -algebra of $B(\mathcal{H})$ (see the remark following Notation 2.10). If we then denote by U_r , $r \in I_M$, a total set of representatives for the irreducible corepresentations of (M, Δ_M) on Hilbert spaces \mathcal{K}_r , we know by [5] that

$$(\mathcal{B}, \alpha) \cong (\bigoplus_{r \in I_M} \mathcal{K}_r \otimes \mathbb{C}^{k_r}, \bigoplus_{r \in I_M} U_r \otimes 1)$$

as a comodule over the Hopf algebra $\mathcal{A} \subseteq M$, where $k_r < \infty$.

Now if β is the projective corepresentation associated with $\mathcal{G}_{12}U_{13}$, then

$$\beta(x) = U_{13}^*(\alpha \otimes \iota)(x)U_{13}, \quad \text{for all } x \in B(\mathcal{H}) \otimes B(\mathcal{K}).$$

Hence if $\tilde{\mathcal{B}}$ is the linear span of the spectral subspaces of β , then as a comodule, we have

$$\tilde{\mathcal{B}} \cong U^c \times \mathcal{B} \times U,$$

where U^c denotes the contragredient of U and where we denote by \times the tensor product of corepresentations/comodules. But this means that the trivial corepresentation appears in \tilde{B} with multiplicity $\sum g_r k_r$, where g_r is the multiplicity of $U_r \subseteq U \times U^c$. Hence the fixed point algebra of β is finite-dimensional, and by the previous Proposition, J will have as its cardinality the dimension of a maximal abelian subalgebra of the fixed point algebra of β . Hence J is finite. \square

The previous proposition leads to the following considerations. Let (N, Δ_N) be a fixed right Galois co-object for a compact Woronowicz algebra (M, Δ_M) . Then we can make a W^* -category \mathcal{D} by considering as objects the (N, Δ_N) -corepresentations which are (isomorphic to) finite direct sums of irreducible (N, Δ_N) -corepresentations, and as morphisms bounded intertwiners. Then if we denote by \mathcal{C} the tensor W^* -category of finite-dimensional (M, Δ_M) -corepresentations, we can make \mathcal{D} into a right \mathcal{C} -module by the natural composition introduced above:

$$\mathcal{D} \times \mathcal{C} \rightarrow \mathcal{D} : (\mathcal{G}, U) \rightarrow \mathcal{G}_{12} U_{13},$$

while the action of morphisms is simply by tensoring. We can then also turn $F_N := \bigoplus_{r \in I_N} \mathbb{Z}$ into a module over the fusion ring $F_M := \bigoplus_{r \in I_M} \mathbb{Z}$ by means of the fusion rules associated with this categorical construction.

But in fact, there is another different way in which to obtain these fusion rules, making use of the theory developed in [28]. In that paper, Wassermann's multiplicity theory for ergodic compact Lie group actions on C^* -and von Neumann algebras is extended to the setting of compact Woronowicz algebras. Although the paper works in the C^* -algebraic realm and uses right coactions, the results also apply in the von Neumann algebra setting and with left actions, and we make the transition without further comment in explaining these ideas.

Let then (M, Δ_M) be a compact Woronowicz algebra with an ergodic coaction α on a von Neumann algebra A . It is well-known that the crossed product

$$M \rtimes A = \{(x \otimes 1)\alpha(y) \mid x \in M, y \in A\}'' \subseteq B(\mathcal{L}^2(M) \otimes \mathcal{L}^2(A))$$

is a von Neumann algebraic direct sum of type I factors. Let I_α be the set of atoms of the center of $M \rtimes A$, and let F_α be the free abelian group generated by I_α . Then one can turn F_α into a right F_M -module by the following procedure. Let $\{p_s \mid s \in I_\alpha\}$ be the set of minimal projections in the center of $M \rtimes A$, and choose for each $s \in I_\alpha$ a minimal projection $e_s \leq p_s$ in $M \rtimes A$. We can equip the corners $e_s(B(\mathcal{L}^2(M)) \otimes A)e_t$ with a left (M, Δ_M) -coaction α_{st} by the formula

$$\alpha_{st}(z) = (\Sigma \otimes 1)(V_{12}^*(\iota \otimes \alpha)(z)V_{12})(\Sigma \otimes 1).$$

For each $r \in I_M$, $s, t \in I_\alpha$, define $M_{st}^{(r)}$ to be the dimension of the set of (M, Δ_M) -intertwiners between the corepresentation U_r associated with r and α_{st} . Then the action of $r \in I_M$ on an element $t \in I_\alpha$ is defined as

$$t \cdot r := \sum_{s \in I_\alpha} M_{ts}^{(r)} \cdot s.$$

Let now (N, Δ_N) be a right Galois co-object for the compact Woronowicz algebra (M, Δ_M) . Choose $r \in I_N$. Then we can apply the above ideas to the left coaction α_r on $B(\mathcal{H}_r)$, where α_r is the coaction associated with the irreducible projective (N, Δ_N) -corepresentation \mathcal{G}_r pertaining to r . *We claim that the resulting right F_M -module is independent of the choice of r , and coincides precisely with the right F_M -module as constructed after Proposition 3.9.* We will briefly indicate how this can be proven.

We first observe that $M \rtimes B(\mathcal{H}_r)$ equals $\mathcal{G}_r^*(\hat{N} \otimes B(\mathcal{H}_r))\mathcal{G}_r$. Indeed, this follows by the characterization of \hat{N} as a fixed point space, by the pentagon identity for \tilde{V} (and the related pentagon identity for

\mathcal{G}_r), by the fact that \mathcal{G}_r implements α_r , and finally by the characterization of $M \rtimes B(\mathcal{H}_r)$ as the set of elements z in $B(\mathcal{L}^2(M)) \bar{\otimes} B(\mathcal{H}_r)$ satisfying

$$V_{12} z_{13} V_{12}^* = (\iota \otimes \alpha_r)(z).$$

Hence we already see that $I_{\alpha_r} = I_N$.

Choose then for each $s \in I_N$ the element

$$e_s := e_{s,00} \otimes e_{r,00} \in \widehat{N} \bar{\otimes} B(\mathcal{H}_r) \cong M \rtimes B(\mathcal{H}_r)$$

as a minimal projection. Then by transporting all structure with the aid of \mathcal{G} , one sees that the corner $e_s(B(\mathcal{L}^2(M)) \bar{\otimes} B(\mathcal{H}_r))e_t$ is isomorphic to $e_{s,00}B(\mathcal{L}^2(N))e_{t,00}$, equipped with the restriction of the coaction Ad_L (which appears in the proof of Proposition 1.10). This may further be simplified to the coaction

$$\alpha_{st} : B(\mathcal{H}_s, \mathcal{H}_t) \rightarrow M \bar{\otimes} B(\mathcal{H}_s, \mathcal{H}_t) : x \rightarrow \mathcal{G}_s^*(1 \otimes x)\mathcal{G}_t.$$

This final coaction may be interpreted as corresponding to the (ordinary) corepresentation ‘ $\mathcal{G}_s^c \times \mathcal{G}_t$ ’. A Frobenius-type argument then shows that this corepresentation contains some U_u with $u \in I_M$ as much times as \mathcal{G}_t is contained in $\mathcal{G}_s \times U_u$. This shows that the two mentioned fusion rules indeed coincide, and ends our sketch of proof.

To end this section, let us come back to comparing the structures of (N, Δ_N) and $(N, \Delta_N^{\text{op}})$ which we started in Lemma 1.18. We begin by introducing a certain antipode on a subspace of N .

Proposition 3.10. *Let (N, Δ_N) be a right Galois co-object for a compact quantum group (M, Δ_M) . Denote by \mathcal{N} the linear span of the $\tilde{V}_{r,ij}$ in N (see Lemma 1.17). Denote by \mathcal{A} the corresponding subspace of M , which coincides with the Hopf *-algebra associated with (M, Δ_M) (see the remark after Notation 2.10). Then the following statements hold.*

1. *The space \mathcal{N} is a right \mathcal{A} -module.*
2. *If we define the anti-linear map*

$$S_N(\cdot)^* : \mathcal{N} \rightarrow \mathcal{N} : \tilde{V}_{r,ij} \rightarrow \tilde{V}_{r,ji},$$

then for all $x \in \mathcal{N}$ and $y \in \mathcal{A}$, we have

$$S_N(xy)^* = S_N(x)^* S_M(y)^*,$$

*where S_M denotes the antipode of the Hopf *-algebra \mathcal{A} .*

3. *For all $r \in I_N$ and $0 \leq i, j < n_r$, we have*

$$\Lambda_N(S_N(\tilde{V}_{r,ij})^*) = \left(\frac{T_{r,j}}{T_{r,i}} \right)^{1/2} J_{\widehat{N}} \Lambda_N(\tilde{V}_{r,ij}),$$

where $J_{\widehat{N}}$ denotes the modular conjugation for \widehat{N} , given by $e_{r,i} \otimes \overline{e_{r,j}} \rightarrow e_{r,j} \otimes \overline{e_{r,i}}$.

Proof. As the $\tilde{V}_{r,ij}$ form a basis of \mathcal{N} , it is easy to see that $S_N(\cdot)^*$ is well-defined. Moreover, using the formulas in Lemma 1.17.1, the third statement follows immediately. As for the first point, this is an immediate consequence of Proposition 3.9.2.

So the only thing left to show is the second statement, which is at least meaningful by the first part of the proposition.

For ω in the predual of \widehat{N}' , denote by $\bar{\omega}$ the normal functional $x \rightarrow \overline{\omega(x^*)}$ on \widehat{N}' . Let us call a normal functional on \widehat{N}' *elementary* when it is of the form $\overline{e_{r,ij}} \rightarrow \omega(e_{r,ji})$ for some ω in the linear span of the $\omega_{s,kl}$ (see Notation 1.14). Then with ω elementary, we immediately obtain the formula

$$S_N((\omega \otimes \iota)(\tilde{V}))^* = (\bar{\omega} \otimes \iota)(\tilde{V}). \quad (6)$$

Choose now normal functionals ω_1 and ω_2 on respectively $B(\mathcal{L}^2(N))$ and $B(\mathcal{L}^2(M))$ which restrict to elementary functionals on respectively \widehat{N}' and \widehat{M}' . Then by the pentagonal identity for \tilde{V} , we have

$$(\omega_1 \otimes \iota)(\tilde{V})(\omega_2 \otimes \iota)(V) = (\tilde{\omega} \otimes \iota)(\tilde{V}),$$

where $\tilde{\omega}$ is the functional

$$x \in B(\mathcal{L}^2(N)) \rightarrow (\omega_1 \otimes \omega_2)(\tilde{V}_{12}^*(1 \otimes x)\tilde{V}_{12}).$$

By the first part of the proposition, the restriction of $\tilde{\omega}$ to \widehat{N}' is again elementary. Combining these statements with equation (6) (and the corresponding one for S_M), we see that $S_N(\cdot)^*$ is indeed right $S_M(\cdot)^*$ -linear. \square

We can now make the connection between the adjoint coactions of (M, Δ_M) on \widehat{N} and $(M, \Delta_M^{\text{op}})$ on \widehat{N}' respectively (see the remark after Lemma 1.18). Let us first recall that any compact Woronowicz algebra (M, Δ_M) is endowed with an involutive anti-comultiplicative anti-* automorphism R_M , given by the formula

$$R_M(x) = J_{\widehat{M}}x^*J_{\widehat{M}} \quad \text{for all } x \in M.$$

More concretely, we have $R_M(W_{r,ij}) = V_{r,ji}$ for all $r \in I_M$ and $0 \leq i, j < m_r$. We will also denote

$$\mathcal{C}_{\widehat{N}} : \widehat{N} \rightarrow \widehat{N}' : x \rightarrow J_{\widehat{N}}x^*J_{\widehat{N}} = \overline{x^*},$$

and use the same notation for its inverse.

Proposition 3.11. *Let (N, Δ_N) be a right Galois co-object for a compact Woronowicz algebra (M, Δ_M) . Let $(N, \Delta_N^{\text{op}})$ be the co-opposite right Galois co-object for $(M, \Delta_M^{\text{op}})$. Then the right adjoint coaction of $(M, \Delta_M^{\text{op}})$ on $(N^{\text{cop}})^\wedge = \widehat{N}'$ is given by*

$$x \rightarrow (\mathcal{C}_N \otimes R_M)Ad_R(\mathcal{C}_N(x)).$$

Proof. Note that the right adjoint coaction on $B(\overline{\mathcal{H}_r})$ is, by its definition and by Lemma 1.18.1, given as

$$x \rightarrow \tilde{V}^*(x \otimes 1)\tilde{V}.$$

Denote now by $J_{\widehat{N}}$ the modular conjugation for \widehat{N} , which we recall is simply the map

$$\mathcal{L}^2(N) \rightarrow \mathcal{L}^2(N) : e_{r,i} \otimes \overline{e_{r,j}} \rightarrow e_{r,j} \otimes \overline{e_{r,i}}.$$

It is then easily seen that the proposition follows one we can prove the following identity:

$$\tilde{V} = (J_{\widehat{N}} \otimes J_{\widehat{N}})(\Sigma \tilde{W}^* \Sigma)(J_{\widehat{N}} \otimes J_{\widehat{M}}). \quad (7)$$

Now piecewise, the identity (7) corresponds to the identities

$$J_{\widehat{N}} \tilde{W}_{r,ji}^* J_{\widehat{M}} = \tilde{V}_{r,ij}, \quad \text{for all } r \in I_N, 0 \leq i, j < n_r.$$

But in [8], it was proven that $J_{\widehat{N}}xJ_{\widehat{M}} \in N$ for $x \in N$ (see the remark just before Lemma 4.3 in that paper). Hence we only have to check if

$$J_{\widehat{N}} \tilde{W}_{r,ji}^* J_{\widehat{M}} \xi_M = \tilde{V}_{r,ij} \xi_M, \text{ for all } r \in I_N, 0 \leq i, j < n_r.$$

This now follows from an easy computation using Lemma 1.17. \square

Remark: It seems nicer to treat the right adjoint $(M, \Delta_M^{\text{op}})$ -coaction on \widehat{N}' as a left (M, Δ_M) -coaction:

$$\text{Ad}_L : \widehat{N}' \rightarrow M \overline{\otimes} \widehat{N}' : x \rightarrow \Sigma \widetilde{V}^*(x \otimes 1) \widetilde{V} \Sigma.$$

These then localize to left adjoint coactions $\text{Ad}_L^{(r)}$ on the $B(\overline{\mathcal{H}}_r)$. Note that the map Ad_L (as well as the map S_N) in fact already appeared in the proof of Proposition 1.10, and that the $\text{Ad}_L^{(r)}$ are nothing but the $[(N, \Delta_N)]$ -corepresentations of (M, Δ_M) associated with the (N, Δ_N) -corepresentations $\Sigma \widetilde{V}_r \Sigma$ from Theorem 3.2.

4 Reflecting a compact Woronowicz algebra across a Galois co-object

In this section, we will consider in the special case of compact Woronowicz algebras a technique which was introduced in [10]. The following theorem was proven in [10], Proposition 2.1 and Theorem 0.7.

Theorem 4.1. *Let (M, Δ_M) be a compact Woronowicz algebra, and (N, Δ_N) a right Galois co-object for (M, Δ_M) . Denote by $P \subseteq B(\mathcal{L}^2(N))$ the von Neumann algebra which is generated by elements of the form xy^* , where $x, y \in N$. Then P can be made into a von Neumann bialgebra, the comultiplication Δ_P being uniquely determined by the fact that*

$$\Delta_P(xy^*) = \Delta_N(x)\Delta_N(y)^* \quad \text{for all } x, y \in N.$$

The von Neumann bialgebra (P, Δ_P) furthermore admits (not necessarily finite) left and right Δ_P -invariant nsf weights (i.e. is a von Neumann algebraic quantum group in the terminology of [20]).

The following theorem gives a concrete formula for the above invariant weights. We will use the notations introduced in the first three sections (see Notation 1.14 and Notation 2.10).

Theorem 4.2. *Let (M, Δ_M) be a compact Woronowicz algebra, (N, Δ_N) a right Galois co-object for (M, Δ_M) , and (P, Δ_P) the von Neumann bialgebra introduced in Theorem 4.1. Then, up to a positive scalar, the left invariant nsf weight φ_P satisfies*

$$\widetilde{V}_{r,ij} \widetilde{V}_{s,kl}^* \in \mathcal{M}_{\varphi_P}$$

with

$$\varphi_P(\widetilde{V}_{r,ij} \widetilde{V}_{s,kl}^*) = \delta_{r,s} \delta_{i,k} \delta_{j,l} \frac{T_{r,j}}{A_{r,j}},$$

while the right invariant nsf weight ψ_P satisfies, again up to a positive scalar,

$$\widetilde{W}_{r,ij}^* \widetilde{W}_{s,kl} \in \mathcal{M}_{\psi_P}$$

and

$$\psi_P(\widetilde{W}_{r,ij}^* \widetilde{W}_{s,kl}) = \delta_{r,s} \delta_{i,k} \delta_{j,l} \frac{T_{r,i}}{A_{r,i}}.$$

Proof. For the proof of the theorem, we have to explain first how the invariant weights φ_P and ψ_P can be obtained. This goes back to Theorem 4.8 of [8].

Denote by $\nabla_{N,M}^{it}$ the following one-parametergroup of unitaries on $\mathcal{L}^2(N)$:

$$\nabla_{N,M}^{it} = \nabla_{\widehat{N}}^{it} J_{\widehat{N}} \delta_{\widehat{N}}^{it} J_{\widehat{N}},$$

where $\nabla_{\widehat{N}}$ is the modular operator associated with the weight $\varphi_{\widehat{N}}$ on \widehat{N} . On basis vectors, this one-parametergroup is concretely given as

$$\nabla_{N,M}^{it} e_{r,i} \otimes \overline{e_{r,j}} = A_{r,j}^{-it} T_{r,j}^{it} T_{r,i}^{-it} e_{r,i} \otimes \overline{e_{r,j}}.$$

We can then implement on N a one-parametergroup $\sigma_t^{N,M}$, determined by the formula

$$\sigma_t^{N,M}(x) = \nabla_{N,M}^{it} x \nabla_M^{-it}, \quad \text{for all } x \in N,$$

where ∇_M is the modular operator on $\mathcal{L}^2(M)$ associated with φ_M . One verifies that this is well-defined by using the commutation relation

$$(\nabla_M^{it} \otimes 1) \widetilde{W}(\nabla_{N,M}^{-it} \otimes 1) = (1 \otimes \nabla_{\widehat{N}}^{-it}) \widetilde{W}(1 \otimes \nabla_{\widehat{N}}^{it} \delta_{\widehat{N}}^{it}),$$

proven in Proposition 3.20 of [8] (we remark that the P -operator introduced there coincides with $\nabla_{\widehat{N}}$, as the modular element of (M, Δ_M) is trivial). This commutation relation also immediately shows that

$$\sigma_t^{N,M}(\widetilde{V}_{r,ij}) = T_{r,i}^{-it} T_{r,j}^{it} A_{r,j}^{-it} \widetilde{V}_{r,ij},$$

using the notation from Lemma 1.17.1.

Now by construction (see the discussion preceding Lemma 4.4 in [8]), all elements $x \in N$ which are analytic with respect to $\sigma_t^{N,M}$ will lie in the space of square integrable elements of φ_P , by the formula

$$\varphi_P(xx^*) = \varphi_M(\sigma_{-i/2}^{N,M}(x)^* \sigma_{-i/2}^{N,M}(x)).$$

By polarity, we get for $x, y \in N$ analytic w.r.t. $\sigma_t^{N,M}$ that

$$\varphi_P(xy^*) = \varphi_M(\sigma_{-i/2}^{N,M}(y)^* \sigma_{-i/2}^{N,M}(x)).$$

Applying this to $x = \widetilde{V}_{r,ij}$ and $y = \widetilde{V}_{s,kl}$, and using the orthogonality relations between the $\widetilde{V}_{r,ij}$, we immediately get the first formula in the statement of the theorem.

For the second formula, we can use the expression $\psi_P = \varphi_P \circ R_P$, where R_P was an involutory anti-automorphism on P determined by the formula

$$R_P(x) = J_{\widehat{N}} x^* J_{\widehat{N}}, \quad \text{for all } x \in P$$

(see Lemma 4.3 in [8]). In fact, the discussion before that lemma states that, for $x, y \in N$, we have $R_P(xy^*) = R_N(y)^* R_N(x)$, where

$$R_N : N \rightarrow N^{\text{op}} : x \rightarrow J_{\widehat{M}} x^* J_{\widehat{N}}.$$

By applying $R_N(\widetilde{V}_{r,ij})^*$ to ξ_M , we find that

$$R_N(\widetilde{V}_{r,ij}) = \widetilde{W}_{r,ji}$$

(see also the proof of Proposition 3.11). Applying then $\varphi_P \circ R_P$ to $\widetilde{W}_{r,ij}^* \widetilde{W}_{s,kl}$ and using the first part of the proof, the expression for ψ_P as in the statement of the theorem follows. □

From the formulas in Theorem 4.2, we can draw the following conclusions.

Proposition 4.3. *Let (M, Δ_M) , (N, Δ_N) and (P, Δ_P) be as in the foregoing theorem.*

1. *If (P, Δ_P) is a compact Woronowicz algebra (that is, if φ_P and ψ_P are finite), then all $n_r < \infty$, i.e. all irreducible (N, Δ_N) -corepresentations of (M, Δ_M) are finite-dimensional.*
2. *Conversely, if one of the irreducible (N, Δ_N) -corepresentations for (M, Δ_M) is finite-dimensional, then they all are, and then (P, Δ_P) is a compact Woronowicz algebra.*
3. *If (P, Δ_P) is unimodular (that is, if φ_P is a multiple of ψ_P), then there exist positive numbers d_r such that $A_r = d_r^2 T_r^2$ (where the A_r were introduced in Notation 2.10).*

Remark: If the condition in the third point is satisfied, then one could interpret d_r as a (finite!) relative quantum dimension of the irreducible (N, Δ_N) -corepresentation corresponding to r , in analogy with the case of ordinary irreducible corepresentations (compare with Proposition 2.2). Here the relativity refers to one irreducible (N, Δ_N) -corepresentation w.r.t. another, as $\delta_{\hat{N}}$ is only determined up to a positive scalar. We note that we do not know of any particular examples where (P, Δ_P) is not unimodular, so it could well be that this condition is *always* fulfilled.

Proof. The third point follows immediately from the formulas in the previous theorem combined with Lemma 1.17, as there then exists a positive number $c > 0$ such that

$$\frac{A_{r,j}}{A_{r,i}} = c \frac{T_{r,j}^2}{T_{r,i}^2}.$$

If (P, Δ_P) is moreover compact, then for any $r \in I_N$ and $0 \leq i < n_r$, we get, by using the unitarity of \widetilde{W}_r (and the normality of ψ_P), that

$$\begin{aligned} \psi_P(1) &= \psi_P\left(\sum_{i=0}^{n_r-1} \widetilde{W}_{r,ij}^* \widetilde{W}_{r,ij}\right) \\ &= \sum_{i=0}^{n_r-1} \psi_P(\widetilde{W}_{r,ij}^* \widetilde{W}_{r,ij}) \\ &= \frac{1}{d_r^2} \sum_{i=0}^{n_r-1} \frac{1}{T_{i,r}} < \infty. \end{aligned}$$

As the $T_{r,i}$ are summable, the final sum must necessarily be finite, i.e. $n_r < \infty$.

Finally, suppose that (M, Δ_M) has a finite-dimensional irreducible (N, Δ_N) -corepresentation, say corresponding to the index value r . Then as $\psi_P(1) = \psi_P(\sum_{i=0}^{n_r-1} \widetilde{W}_{r,ij}^* \widetilde{W}_{r,ij})$, we see that ψ_P is finite, and hence (P, Δ_P) is a compact Woronowicz algebra. By the second point, also all other irreducible (N, Δ_N) -corepresentations of (M, Δ_M) are finite-dimensional. □

Remark: In case the irreducible (N, Δ_N) -corepresentations are finite-dimensional, the linear span of the $\widetilde{W}_{r,ij}^*$ generates inside N a purely algebraic Galois co-object \mathcal{N} for the Hopf algebra \mathcal{A} inside (M, Δ_M) . Conversely, if one starts with a Galois co-object for \mathcal{A} , satisfying some suitable relations with the *-structure, we can in essence develop the whole theory so far in an algebraic way, and then *necessarily* the reflection will correspond to a compact quantum group (this was essentially already observed in [7], see also [32]). As it turns out, there *do* exist interesting Galois co-objects which are of a non-algebraic type (see the final section), which was part of the motivation for writing this paper.

5 Galois co-objects and projective corepresentations for compact Kac algebras

In this short section, we show that when one deals with compact Kac algebras (see Definition 1.1), one is *always* in the algebraic setup (see the remark at the end of the previous section).

Proposition 5.1. *Let (N, Δ_N) be a Galois co-object for a compact Kac algebra (M, Δ_M) , and let (P, Δ_P) be the reflected von Neumann bialgebra as obtained in Theorem 4.1. Then also (P, Δ_P) is a compact Kac algebra.*

Proof. As we recalled in Theorem 2.9, the modular element $\delta_{\widehat{N}}^{it}$ satisfies $\alpha_{\widehat{N}}(\delta_{\widehat{N}}^{it}) = \delta_{\widehat{N}}^{it} \otimes \delta_{\widehat{M}}^{it}$. However, for a compact Kac algebra, $\delta_{\widehat{M}} = 1$. By ergodicity of $\alpha_{\widehat{N}}$, we then conclude that we can take $\delta_{\widehat{N}} = 1$.

From Theorem 4.2 and the normality of ψ_P , we then find

$$\begin{aligned} \psi_P(1) &= \psi_P\left(\sum_{i=0}^{n_r-1} \widetilde{W}_{r,ij}^* \widetilde{W}_{r,ij}\right) \\ &= \sum_{i=0}^{n_r-1} \psi_P(\widetilde{W}_{r,ij}^* \widetilde{W}_{r,ij}) \\ &= \sum_{i=0}^{n_r-1} T_{i,r} = 1, \end{aligned}$$

so that ψ_P is finite, and (P, Δ_P) thus a compact Woronowicz algebra.

But then (P, Δ_P) is in particular unimodular, so that the third point of Proposition 4.3 gives us that T_r is a scalar matrix, and hence just $\frac{1}{n_r}$ times the unit matrix on \mathcal{H}_r . This shows that the one-parameter group $\nabla_{N,M}^{it}$, which we introduced in the course of the proof of Theorem 4.2, is trivial. As $\sigma_t^{\varphi_P}(xy^*) = \sigma_t^{N,M}(x)\sigma_t^{N,M}(y)^*$ for all $x, y \in N$ (which follows from the fact that $\sigma_t^{N,M}$ is actually the restriction to N of the modular automorphism group of the balanced weight $\varphi_P \oplus \tau_M$ on $\begin{pmatrix} P & N \\ N^{\text{op}} & M \end{pmatrix}$), we get that $\sigma_t^{\varphi_P}$ is trivial, and hence φ_P is a trace. This concludes the proof. \square

Combining the previous proposition with Theorem 3.2 and Proposition 4.3, we obtain the following corollary.

Corollary 5.2. *Let (M, Δ_M) be a compact Kac algebra. If \mathcal{H} is a Hilbert space, and $\alpha : B(\mathcal{H}) \rightarrow M \overline{\otimes} B(\mathcal{H})$ a coaction, then the following statements hold.*

1. *The trace on $B(\mathcal{H})$ is α -invariant.*
2. *If α is ergodic, then \mathcal{H} is finite-dimensional.*

In particular, this says that the invariant state associated with an ergodic coaction of a compact Kac algebra on a type I-factor is tracial. Note that this is *not* true for ergodic coactions of Kac algebras on *arbitrary* von Neumann algebras (counterexamples can be found in [4]). It would be nice to have a more direct proof of the above corollary, but we were not able to produce one.

6 Projective corepresentations of finite-dimensional Kac algebras

In this section, we will briefly review what can be said concerning the situation of finite-dimensional compact Woronowicz (and hence Kac) algebras.

Proposition 6.1. *Let (M, Δ_M) be a finite-dimensional Kac algebra, and (N, Δ_N) a right Galois co-object for (M, Δ_M) . Then N is finite-dimensional with $\dim(N) = \dim(M)$, and (N, Δ_N) is cleft.*

Proof. Choose $r \in I_N$. Then as the right coaction $\text{Ad}_R^{(r)}$ of (M, Δ_M) on $B(\mathcal{H}_r)$ is ergodic, it is easy to see *immediately* that $n_r = \dim(\mathcal{H}_r)$ is finite. Then take $r \in I_N$ fixed. As N is the σ -weak closure of the set $\{\widetilde{W}_{r,ij}^* m \mid 0 \leq i, j \leq n_r, m \in M\}$ by Lemma 2.5, we see that N is finite-dimensional. As \widetilde{W}^* gives a unitary from $\mathcal{L}^2(N) \otimes \mathcal{L}^2(M)$ to $\mathcal{L}^2(N) \otimes \mathcal{L}^2(N)$, necessarily $\dim(N) = \dim(M)$.

Now disregarding the $*$ -structures, we get that (N, Δ_N) is a Galois co-object for the Hopf algebra (M, Δ_M) . It is then well-known that (N, Δ_N) is cleft in this weaker form (see for example the remark following Corollary 3.2.4 in [26]). But this means that $N \cong M$ as right M -modules. As M is a direct sum of matrix algebras, it is easy to see that we can in fact find a unitary $u : \mathcal{L}^2(N) \rightarrow \mathcal{L}^2(M)$ such that $uN = M$. Hence we may identify $\mathcal{L}^2(N)$ with $\mathcal{L}^2(M)$ and N with M . We can then consider $\Omega = \Delta_N(1_M)$. By right linearity of Δ_N , we then get $\Delta_N(x) = \Omega \Delta_M(x)$ for all $x \in N$, and by coassociativity of Δ_N we have that Ω satisfies the 2-cocycle relation. Hence (N, Δ_N) is cleft. \square

Remark: Finite Galois co-objects (in the operator algebra context) have also been dealt with in the papers [12], [31] and, in a more general setting, [17].

For later purposes, we also introduce the following definition of a *non-degenerate* 2-cocycle (see Definition 1.19 for the general notion of a unitary 2-cocycle).

Definition 6.2. *Let (M, Δ_M) be a finite-dimensional compact Kac algebra, and Ω a unitary 2-cocycle for (M, Δ_M) . We call Ω non-degenerate if the associated Galois object \widehat{N} is a (finite-dimensional) type I-factor.*

This terminology was introduced in [1]. One observes that, as $(\widehat{N}, \alpha_{\widehat{N}})$ is then in fact also a projective (right) corepresentation for $(\widehat{M}, \Delta_{\widehat{M}})$, we can create from it a Galois co-object for $(\widehat{M}, \Delta_{\widehat{M}})$, which will then necessarily also be cleft. If we denote by $\widehat{\Omega} \in \widehat{M} \otimes \widehat{M}$ an implementing unitary 2-cocycle, then $\widehat{\Omega}$ turns out to be non-degenerate again, and the correspondence $[\Omega] \rightarrow [\widehat{\Omega}]$ between cohomology classes of non-degenerate 2-cocycles of resp. (M, Δ_M) and $(\widehat{M}, \Delta_{\widehat{M}})$ is a bijection. (For the proof of this result, we refer again to [1].) To give a simple example, consider a finite abelian group G . Then the bicharacter on $\widehat{G} \times G$ given by evaluation gives a non-degenerate 2-cocycle function Ω on $G \times \widehat{G}$ by the formula $((g, \chi), (h, \chi')) \rightarrow \chi(h)$, and its dual is simply the same construction applied to the evaluation bicharacter on $\widehat{G} \times G$.

7 Projective corepresentations for cocommutative Kac algebras

As a second special case, we will consider compact Woronowicz algebras (M, Δ_M) which have a cocommutative coproduct: $\Delta_M^{\text{op}} = \Delta_M$. It is not so difficult to show that (M, Δ_M) is then Kac, and

in fact that $M = \mathcal{L}(\Gamma)$ for some (countable) discrete group Γ , the coproduct being given by

$$\Delta_M(\lambda_g) = \lambda_g \otimes \lambda_g \quad \text{for all } g \in \Gamma,$$

where the λ_g denote the standard unitary generators in the group von Neumann algebra $\mathcal{L}(\Gamma)$. We will in the following denote $\mathcal{L}(\Gamma)$ as shorthand for $(\mathcal{L}(\Gamma), \Delta_{\mathcal{L}(\Gamma)})$, and we denote the invariant trace by τ . The dual discrete Woronowicz algebra $(\widehat{M}, \Delta_{\widehat{M}})$ is then simply the function space $l^\infty(\Gamma)$, equipped with the coproduct

$$\Delta_{\widehat{M}} : l^\infty(\Gamma) \rightarrow l^\infty(\Gamma) \bar{\otimes} l^\infty(\Gamma) \cong l^\infty(\Gamma \times \Gamma)$$

such that

$$\Delta_{\widehat{M}}(f)(g, h) = f(gh) \quad \text{for all } g, h \in \Gamma.$$

We will in the following write $\mathcal{L}^2(\mathcal{L}(\Gamma)) = l^2(\Gamma)$ of course, and then, with δ_g being the Dirac function at the point $g \in \Gamma$, we have

$$\Lambda_{\mathcal{L}(\Gamma)}(\lambda_g) = \delta_g.$$

As group von Neumann algebras are in particular Kac algebras, we know from Corollary 5.2 that they can only act ergodically on type I factors which are of finite type. Let us give a more immediate proof of this fact in this particular case.

Lemma 7.1. *Let Γ be a discrete group. Let B be a von Neumann algebra, and suppose that we have given an ergodic coaction*

$$\alpha : B \rightarrow \mathcal{L}(\Gamma) \otimes B.$$

If we denote by ϕ_B the unique α -invariant state on B , then ϕ_B is a trace.

Proof. For each $g \in \Gamma$, denote $B_g = \{x \in B \mid \alpha(x) = \lambda_g \otimes x\}$. As α is ergodic, it is easily seen that each B_g is either zero- or one-dimensional. It is further immediate that $B_g \cdot B_h \subseteq B_{gh}$ for all $g, h \in \Gamma$, and that $B_g^* = B_{g^{-1}}$. Therefore, whenever B_g is not zero-dimensional, we may assume that $B_g = \mathbb{C}u_g$ with u_g a unitary. We may moreover assume that $u_g^* = u_{g^{-1}}$. When $B_g = 0$, we will denote $u_g = 0$.

We claim that the linear span of the u_g is σ -weakly dense in B . Indeed, if this was not the case, then we could find a nonzero $x \in B$ with $\phi_B(xu_g) = 0$ for all $g \in \Gamma$. But as $\phi_B = (\tau \otimes \iota)\alpha$ by definition, this would imply that

$$(\tau \otimes \iota)(\alpha(x)(\lambda_g \otimes 1))u_g = 0 \quad \text{for all } g \in \Gamma.$$

Now an easy computation shows that for all $g \in \Gamma$, we have

$$(\tau \otimes \iota)(\alpha(x)(\lambda_g \otimes 1)) \in B_g^*.$$

Hence we in fact have

$$(\tau \otimes \iota)(\alpha(x)(\lambda_g \otimes 1)) = 0 \quad \text{for all } g \in \Gamma.$$

This implies $\alpha(x) = 0$, and so $x = 0$, which is a contradiction.

It is now enough to prove that $\phi_B(u_g u_h) = \phi_B(u_h u_g)$ for all $g, h \in \Gamma$. But the left hand side is a multiple of $\phi_B(u_{gh})$, which is zero in case $g \neq h^{-1}$. Similarly, the right hand side is zero in case $g \neq h^{-1}$. As both sides equal 1 when $g = h^{-1}$, we are done. \square

Remark: General coactions of group von Neumann algebras (or rather, of the associated group C^* -algebras), have been studied in the theory of *Fell bundles* over groups (see for example [25]). The intuitive connection between these notions is essentially contained the above proof.

Corollary 7.2. *Let Γ be a discrete group, \mathcal{H} a Hilbert space, and*

$$\alpha : B(\mathcal{H}) \rightarrow \mathcal{L}(\Gamma) \bar{\otimes} B(\mathcal{H})$$

an ergodic coaction of $\mathcal{L}(\Gamma)$ on $B(\mathcal{H})$. Then the following statements hold.

1. *The dimension of \mathcal{H} is finite.*
2. *There exists a finite subgroup H of Γ such that*

$$\alpha(B(\mathcal{H})) \subseteq \mathcal{L}(H) \otimes B(\mathcal{H}),$$

and such that, denoting by β the coaction α with range restricted to $\mathcal{L}(H) \otimes B(\mathcal{H})$, the couple $(B(\mathcal{H}), \beta)$ is a (left) Galois object for $\mathcal{L}(H)$.

Remark: The notion of a Galois object was introduced in the second section. In the finite-dimensional setting, it may be defined as follows: let A be a finite-dimensional Hopf $*$ -algebra with a left coaction β on a finite-dimensional $*$ -algebra B . Then (B, β) is called a left Galois object for A if the map

$$B \otimes B \rightarrow A \otimes B : x \otimes y \rightarrow \beta(x)(1 \otimes y)$$

is a bijection.

Proof. By the previous lemma, we know that $B(\mathcal{H})$ admits a tracial state. Hence \mathcal{H} must be finite.

As for the second point, this is rather a corollary of the proof of the previous proposition. For, using the notation introduced there, denote by H the set of elements g in Γ for which $u_g \neq 0$. As the u_g are orthogonal to each other with respect to the α -invariant state, we must have that H is finite. As $u_g \cdot u_h$ is a non-zero multiple of u_{gh} , and $u_g^* = u_{g^{-1}}$, we must have that H is a finite group. It is then immediate that indeed $\alpha(B(\mathcal{H})) \subseteq \mathcal{L}(H) \otimes B(\mathcal{H})$.

The coaction β of $\mathcal{L}(H)$ on $B(\mathcal{H})$ is then clearly also an ergodic action, with the ordinary (normalized) trace tr as its invariant state. This implies that the map

$$\mathcal{L}^2(B(\mathcal{H}), \text{tr}) \otimes \mathcal{L}^2(B(\mathcal{H}), \text{tr}) \rightarrow (\mathcal{L}^2(\mathcal{L}(H)), \tau) \otimes \mathcal{L}^2(B(\mathcal{H}), \text{tr}) : x \otimes y \rightarrow \beta(x)(1 \otimes y)$$

is isometric and thus injective. As the u_g with $g \in H$ form an orthonormal basis of $B(\mathcal{H})$, we have that the order of H equals the square of the dimension of \mathcal{H} . Hence a comparison of dimensions shows that the above map is also surjective, which proves that $(B(\mathcal{H}), \beta)$ is a left Galois object for $\mathcal{L}(H)$. □

Proposition 7.3. *Let Γ be a discrete group, and let (N, Δ_N) be a right Galois co-object for $\mathcal{L}(\Gamma)$. Then there exists a finite subgroup $H \subseteq \Gamma$ and a non-degenerate 2-cocycle Ω for $\mathcal{L}(H)$, such that (N, Δ_N) is isomorphic to the cleft Galois co-object induced by Ω (considered as a 2-cocycle for $\mathcal{L}(\Gamma)$).*

Proof. Let (N, Δ_N) be a right Galois co-object for $\mathcal{L}(\Gamma)$. Using the terminology introduced in Definition 3.6, choose an irreducible $[(N, \Delta_N)]$ -corepresentation

$$\alpha : B(\mathcal{H}) \rightarrow \mathcal{L}(\Gamma) \otimes B(\mathcal{H})$$

of $\mathcal{L}(\Gamma)$ on a Hilbert space \mathcal{H} (for example, one of the $\text{Ad}_L^{(r)}$, see the end of the third section). By the previous proposition, we know that \mathcal{H} is finite-dimensional, and that we can choose a minimal finite subgroup $H \subseteq \Gamma$ for which

$$\alpha(B(\mathcal{H})) \subseteq \mathcal{L}(H) \otimes B(\mathcal{H}).$$

We moreover know that the corresponding coaction β of $\mathcal{L}(H)$ on $B(\mathcal{H})$ is then a Galois object. This means that the Galois co-object (N_H, Δ_{N_H}) which is associated with β (as a projective corepresentation) may be taken to be equal to $(\mathcal{L}(H), \Omega \Delta_{\mathcal{L}(H)}(\cdot))$, where $\Omega \in \mathcal{L}(H) \otimes \mathcal{L}(H)$ is a *non-degenerate* unitary 2-cocycle (see the final remarks of the previous section). Denote further $(\tilde{N}, \Delta_{\tilde{N}}) := (\mathcal{L}(\Gamma), \Omega \Delta_{\mathcal{L}(\Gamma)}(\cdot))$, which is a cleft Galois co-object for $\mathcal{L}(\Gamma)$.

Let then \mathcal{G} be a projective (N_H, Δ_{N_H}) -corepresentation which implements β . As $N_H = \mathcal{L}(H) \subseteq \tilde{N} = \mathcal{L}(\Gamma)$, we may interpret \mathcal{G} to be an element of $\tilde{N} \otimes B(\mathcal{H})$. It is trivial to see that \mathcal{G} is then an $(\tilde{N}, \Delta_{\tilde{N}})$ -corepresentation which implements α . Therefore (N, Δ_N) is isomorphic to $(\tilde{N}, \Delta_{\tilde{N}})$ as a right Galois co-object for $\mathcal{L}(\Gamma)$ (see the remark after Definition 3.6), which proves the proposition. \square

Corollary 7.4. *If Γ is a torsionless discrete group, or more generally, a group with no finite subgroups of square order, then any Galois co-object for $(\mathcal{L}(\Gamma), \Delta_{\mathcal{L}(\Gamma)})$ is isomorphic to $(\mathcal{L}(\Gamma), \Delta_{\mathcal{L}(\Gamma)})$ as a right Galois co-object.*

In particular, any unitary 2-cocycle for $\mathcal{L}(\Gamma)$ is then a 2-coboundary (see the remark after Example 1.20).

Proof. The statement concerning torsionless discrete groups is of course an immediate consequence of the previous proposition. As for the statement concerning the case when there are no finite subgroups of square order, observe that if K is any finite group for which some $B(\mathcal{H})$ allows a Galois object structure for $l^\infty(K)$, then necessarily $|K| = \dim(H)^2$. \square

Remarks:

1. For finite groups, Proposition 7.3 was proven in [24] (see also [13]).
2. In [16], it is shown that for *any* group Γ , all *quasi-symmetric* 2-cocycles for $\mathcal{L}(\Gamma)$, i.e. cocycles which also satisfy $\Sigma \Omega \Sigma = \eta \Omega$ for some $\eta \in S^1$, are coboundaries. (The authors weaken this to allow unitaries satisfying the 2-cocycle condition up to a scalar, but it is possible to show that, in any compact Woronowicz algebra, such unitaries are *automatically* 2-cocycles).
3. One can also easily describe the Galois objects dual to the Galois co-objects appearing in Proposition 7.3. Namely, if we have given a discrete group Γ , a finite subgroup H and a non-degenerate 2-cocycle Ω for $\mathcal{L}(H)$, let $\gamma : B(\mathcal{H}) \rightarrow B(\mathcal{H}) \otimes l^\infty(H)$ be the Galois object for $l^\infty(H)$ dual to the Galois co-object associated with Ω . Then the dual of the Galois co-object for $\mathcal{L}(\Gamma)$ associated with (H, Ω) is the *induction* of γ to Γ . The underlying von Neumann algebra of this construction consists of the set of all elements $x \in B(\mathcal{H}) \otimes l^\infty(\Gamma)$ for which

$$(\gamma \otimes \iota)(x) = (\iota \otimes \beta_{l^\infty(H)})(x),$$

where $\beta_{l^\infty(H)}$ is the coaction associated with the left translation action of H on Γ . The right coaction α of $l^\infty(\Gamma)$ on this von Neumann algebra is then simply given by right translation, i.e. $\alpha(x) := (\iota \otimes \Delta_{l^\infty(\Gamma)})(x)$.

The projective corepresentations associated with the Galois co-objects for $\mathcal{L}(\Gamma)$ can be determined as follows.

Proposition 7.5. *Let Γ be a discrete group with a finite subgroup H . Let Ω be a non-degenerate 2-cocycle for $\mathcal{L}(H)$, and let $\mathcal{G} \in \mathcal{L}(H) \otimes B(\mathcal{H})$ be the associated irreducible Ω -corepresentation on some Hilbert space \mathcal{H} .*

Then with (N, Δ_N) the cleft Galois co-object for $\mathcal{L}(\Gamma)$ associated with $\Omega \in \mathcal{L}(\Gamma) \overline{\otimes} \mathcal{L}(\Gamma)$, we have $I_N \cong H \setminus \Gamma$, and a maximal set of irreducible non-isomorphic (N, Δ_N) -corepresentations is given by the set

$$\mathcal{G}_{Hg} := \mathcal{G}(\lambda_{s(Hg)} \otimes 1) \in \mathcal{L}(\Gamma) \otimes B(\mathcal{H}),$$

where $s : H \setminus \Gamma \rightarrow \Gamma$ is a fixed section for $\Gamma \rightarrow H \setminus \Gamma$.

Remark: As Ω is assumed to be non-degenerate for $\mathcal{L}(H)$, the unitary \mathcal{G} is indeed the *unique* Ω -corepresentation for $\mathcal{L}(H)$, up to isomorphism.

Proof. It is immediately seen that the right regular (N, Δ_N) -corepresentation for $\mathcal{L}(\Gamma)$ equals $\tilde{V} = \Omega V$, while the left regular $(N^{\text{op}}, \Delta_{N^{\text{op}}})$ -corepresentation equals $\tilde{W} = \Omega_H^* W$. For $g \in \Gamma$, let $\delta_{Hg} \in B(l^2(\Gamma))$ be the indicator function for the coset Hg . Clearly, δ_{Hg} commutes pointwise with $\mathcal{L}(H)$. Using then the definition of \hat{N} as a fixed point set (see Proposition 1.10), it is easy to check that $\delta_{Hg} \in \hat{N}$. Using the second definition of \hat{N} as the closure of the right leg of \tilde{W} (see again Proposition 1.10), we get that in fact $\delta_{Hg} \in \mathcal{Z}(\hat{N})$, the center of \hat{N} .

We claim now that $I_N \cong H \setminus \Gamma$, and that the $\tilde{V}_{Hg} := \tilde{V}(\delta_{Hg} \otimes 1)$ are the irreducible components of \tilde{V} . Indeed, as $\tilde{V} = \bigoplus_{g \in H} \tilde{V}_{Hg}$, it is enough to show that each of the sets $\{(\iota \otimes \omega)(\tilde{V}_{Hg}) \mid \omega \in \mathcal{L}(\Gamma)_*\}$ is a type I -factor. But denoting $V_H = \sum_{h \in H} \delta_h \otimes \lambda_h$, an easy computation shows that

$$\tilde{V}_{Hg} = (\rho_g \otimes 1) \Omega V_H (\rho_g^* \otimes \lambda_g),$$

where ρ_g is a right translation operator on $l^2(\Gamma)$, i.e. $\rho_g \delta_k = \delta_{kg}$ for $g, k \in \Gamma$. As we assumed that Ω is a non-degenerate 2-cocycle, we know that $\{(\iota \otimes \omega)(\Omega V_H) \mid \omega \in \mathcal{L}(\Gamma)_*\}$ is a type I -factor. This proves the claim.

Now the irreducible Ω -corepresentation $\Sigma \tilde{V}_{Hg} \Sigma$ of $\mathcal{L}(\Gamma)$ is immediately seen to be isomorphic to the Ω -corepresentation \mathcal{G}_{gH} in the statement of the proposition, while \tilde{V}_H is isomorphic to \mathcal{G} as an Ω -corepresentation for $\mathcal{L}(H)$. This then finishes the proof. \square

Remarks:

1. With the help of this proposition, one can show that if H and K are two finite subgroups of Γ , with respective non-degenerate 2-cocycles Ω_H and Ω_K , then the associated Galois co-objects (N, Δ_N) and $(\tilde{N}, \Delta_{\tilde{N}})$ for $\mathcal{L}(\Gamma)$ are isomorphic iff there exists $g \in \Gamma$ and a unitary $u \in \mathcal{L}(H)$ with $g^{-1}Hg = K$ and

$$(\lambda_g \otimes \lambda_g) \Omega_2 (\lambda_g^* \otimes \lambda_g^*) = (u^* \otimes u^*) \Omega_1 \Delta_{\mathcal{L}(H)}(u).$$

2. One can also be more specific on when the projective corepresentations associated with two given (N, Δ_N) -corepresentations as above are actually isomorphic. Namely, using the notation as in the statement of the proposition, let α_g be the coaction of $\mathcal{L}(\Gamma)$ on $B(\mathcal{H})$ implemented by \mathcal{G}_{Hg} . We may assume that α_e is the ‘extension’ of the coaction β of $\mathcal{L}(H)$ on $B(\mathcal{H})$ implemented by \mathcal{G} . Then we have $\alpha_g \cong \alpha_e$ iff $gHg^{-1} = H$ and Ω is coboundary equivalent to $(\lambda_g \otimes \lambda_g) \Omega (\lambda_g^* \otimes \lambda_g^*)$ (inside $\mathcal{L}(H)$).

8 A projective representation for $SU_q(2)$

In this section, we want to consider one special and non-trivial example of a projective representation of the compact quantum group $SU_q(2)$. This projective representation will be nothing else but (a completion of) its action on the standard Podleś sphere.

Let us first recall the definition of $SU_q(2)$ on the von Neumann algebra level. *For the rest of this section, we fix a number $0 < q < 1$.*

Definition 8.1. Denote $\mathcal{L}^2(SU_q(2)) = l^2(\mathbb{N}) \otimes \overline{l^2(\mathbb{N})} \otimes l^2(\mathbb{Z})$. Consider on it the operators

$$a = \sum_{k \in \mathbb{N}_0} \sqrt{1 - q^{2k}} e_{k-1,k} \otimes 1 \otimes 1,$$

$$b = \left(\sum_{k \in \mathbb{N}} q^k e_{kk} \right) \otimes 1 \otimes S,$$

where S denotes the forward bilateral shift.

Then the compact Woronowicz algebra $(\mathcal{L}^\infty(SU_q(2)), \Delta_+)$ consists of the von Neumann algebra

$$\mathcal{L}^\infty(SU_q(2)) = B(l^2(I_+)) \bar{\otimes} 1 \bar{\otimes} \mathcal{L}(\mathbb{Z}) \subseteq B(\mathcal{L}^2(SU_q(2))),$$

equipped with the unique unital normal $*$ -homomorphism

$$\Delta_+ : \mathcal{L}^\infty(SU_q(2)) \rightarrow \mathcal{L}^\infty(SU_q(2)) \bar{\otimes} \mathcal{L}^\infty(SU_q(2))$$

which satisfies

$$\begin{cases} \Delta_+(a) = a \otimes a - qb^* \otimes b \\ \Delta_+(b) = b \otimes a + a^* \otimes b. \end{cases}$$

Its invariant state φ_+ is given by the formula

$$\varphi_+(e_{ij} \otimes 1 \otimes S^k) = \delta_{i,j} \delta_{k,0} (1 - q^2) q^{2k}, \quad \text{for all } i, j \in \mathbb{N}, k \in \mathbb{Z},$$

and we may identify $\mathcal{L}^2(SU_q(2))$ with the Hilbert space in the GNS-construction for φ_+ by putting $\xi_M = \sqrt{1 - q^2} \sum_{i \in \mathbb{N}} q^i e_i \otimes \bar{e}_i \otimes e_0$.

The definition of the standard Podleś sphere and the associated action of $SU_q(2)$ takes the following form on the von Neumann algebraic level.

Definition 8.2. Define $\mathcal{L}^\infty(S_{q_0}^2)$ to be the von Neumann algebra inside $\mathcal{L}^\infty(SU_q(2))$ generated by $X = qb^*a$ and $Z = b^*b$. Then Δ_+ restricts to a left (ergodic) coaction α of $\mathcal{L}^\infty(SU_q(2))$ on $\mathcal{L}^\infty(S_{q_0}^2)$. We say that this coaction corresponds to ‘the action of $SU_q(2)$ on the standard Podleś sphere’.

One can show that $\mathcal{L}^\infty(S_{q_0}^2)$ may be identified with the von Neumann algebra $B(l^2(\mathbb{N}))$, in such a way that

$$X \rightarrow \sum_{k \in \mathbb{N}_0} q^k \sqrt{1 - q^{2k}} e_{k-1,k}$$

$$Z \rightarrow \sum_{k \in \mathbb{N}} q^{2k} e_{kk}.$$

Under this correspondence, the α -invariant state on $\mathcal{L}^\infty(S_{q_0}^2) = B(l^2(\mathbb{N}))$ equals

$$\phi_\alpha(e_{ij}) = \delta_{ij} (1 - q^2) q^{2i}.$$

In the terminology of the present paper, the coaction α is thus an irreducible projective representation of $SU_q(2)$ on an infinite dimensional Hilbert space. In [9], we computed the associated Galois co-object. To introduce it, let us first recall some notations from q -analysis (see [14]).

Notation 8.3. For $n \in \mathbb{N} \cup \{\infty\}$ and $a \in \mathbb{C}$, we denote

$$(a; q)_n = \prod_{k=0}^{n-1} (1 - q^k a),$$

which determines analytic functions in the variable a with no zeroes in the open unit disc.

For $n \in \mathbb{N}$ and $a \in \mathbb{C}$, we denote by $p_n(x; a, 0 \mid q)$ the Wall polynomial of degree n with parameter value a ; so

$$p_n(x; a, 0 \mid q) = {}_2\varphi_1(q^{-n}, 0; qa \mid q, qx),$$

where ${}_2\varphi_1$ denotes Heine's basic hypergeometric function.

Proposition 8.4. Let $\mathcal{L}^2(N) = l^2(\mathbb{Z}) \otimes \overline{l^2(\mathbb{Z})} \otimes l^2(\mathbb{Z})$. Denote by v the operator $S^* \otimes 1 \otimes 1$, where S denotes the forward bilateral shift, and by L_{0+} the operator such that

$$L_{0+}(e_n \otimes \bar{e}_m \otimes e_k) = (q^{2n+2}; q^2)_\infty^{1/2} e_n \otimes \bar{e}_m \otimes e_k,$$

so $L_{0+} = u(q^2 b^* b; q^2)_\infty^{1/2}$ with u the canonical isometric inclusion of $l^2(\mathbb{N}) \otimes \overline{l^2(\mathbb{N})} \otimes l^2(\mathbb{Z})$ into $l^2(\mathbb{Z}) \otimes \overline{l^2(\mathbb{Z})} \otimes l^2(\mathbb{Z})$. Denote N for the σ -weak closure of the right $\mathcal{L}^\infty(SU_q(2))$ -module generated by the elements $v^n L_{0+}$, where $n \in \mathbb{Z}$. (One easily shows that N equals the set $B(l^2(\mathbb{N}), l^2(\mathbb{Z})) \bar{\otimes} 1 \otimes \mathcal{L}(\mathbb{Z})$.)

Then there exists a unique Galois co-object structure (N, Δ_N) on N for which

$$\Delta_N(v^n L_{0+}) = (v^n \otimes v^n) \cdot \left(\sum_{p=0}^{\infty} (q^2; q^2)_p^{-1} v^p L_{0+} b^p \otimes v^p L_{0+} (-qb^*)^p \right),$$

the right hand side converging in norm.

The coaction α of $\mathcal{L}^\infty(SU_q(2))$ on $\mathcal{L}^\infty(S_{q0}^2)$ is then an irreducible $[(N, \Delta_N)]$ -corepresentation, and an associated implementing (N, Δ_N) -corepresentation \mathcal{G} is determined by the following formula: denoting $\mathcal{G} = \sum_{s,t \in \mathbb{N}} \mathcal{G}_{ts} \otimes e_{ts} \in N \bar{\otimes} B(l^2(\mathbb{N}))$, we have, for $0 \leq t \leq s$, that

$$\mathcal{G}_{ts} = q^{t(t-s)} \left(\frac{(q^2; q^2)_s}{(q^2; q^2)_t} \right)^{1/2} (q^2; q^2)_{s-t}^{-1} \cdot v^{s+t} L_{0+} b^{s-t} \cdot p_t(b^* b; q^{2(s-t)}, 0 \mid q^2),$$

while for $0 \leq s \leq t$, we have

$$\mathcal{G}_{ts} = q^{s(s-t)} \left(\frac{(q^2; q^2)_t}{(q^2; q^2)_s} \right)^{1/2} (q^2; q^2)_{t-s}^{-1} \cdot v^{t+s} L_{0+} (-qb^*)^{t-s} \cdot p_s(b^* b; q^{2(t-s)}, 0 \mid q^2).$$

For the rest of this section, we take (M, Δ_M) to be $(\mathcal{L}^\infty(SU_q(2)), \Delta_{\mathcal{L}^\infty(SU_q(2))})$, and we fix the right Galois co-object (N, Δ_N) for (M, Δ_M) introduced above. We then also keep using the notations introduced above, as well as those from the first four sections.

Our aim now is to describe in more detail the structure of the Galois co-object (N, Δ_N) . In particular, we want to find a complete set of irreducible (N, Δ_N) -corepresentations. This is in fact not so difficult.

Proposition 8.5. Let (N, Δ_N) be the Galois co-object and \mathcal{G} the (N, Δ_N) -corepresentation introduced in Proposition 8.4.

Then the set of unitaries

$$\mathcal{G}^{(n)} := (v^n \otimes 1)\mathcal{G}, \quad n \in \mathbb{Z}$$

forms a complete set of irreducible (N, Δ_N) -corepresentations for $\mathcal{L}^\infty(SU_q(2))$.

In particular, the set I_N of isomorphism classes of irreducible (N, Δ_N) -corepresentations can be naturally identified with \mathbb{Z} .

Proof. It is trivial to see that the $\mathcal{G}^{(n)}$ are indeed irreducible (N, Δ_N) -corepresentations, by the group-like property of v . We then only need to show that the $\mathcal{G}^{(n)}$ are mutually non-isomorphic and have σ -weakly dense linear span in N .

We first prove that all $\mathcal{G}^{(n)}$ are mutually non-isomorphic. As an isomorphism between $\mathcal{G}^{(n)}$ and $\mathcal{G}^{(m)}$ would induce an isomorphism between $\mathcal{G}^{(0)}$ and $\mathcal{G}^{(m-n)}$, it is sufficient to show that $\mathcal{G} = \mathcal{G}^{(0)}$ is not isomorphic to $\mathcal{G}^{(n)}$ for $n \neq 0$. But for this, it is in turn sufficient to show that $L_{0+} = \mathcal{G}_{00}$ is orthogonal to all $\mathcal{G}_{ts}^{(n)}$, by the orthogonality relations in Lemma 1.17.2 (and Theorem 3.2.2).

Now we remark that φ_+ satisfies the property that $\varphi_+(a^m b^k (b^*)^l) = 0$ whenever $m \neq 0$ and $k \neq l$, and likewise with a replaced by a^* . Moreover, one easily computes that the commutation relation $v^* L_{0+} = L_{0+} a^*$ holds. Using then the concrete form for the \mathcal{G}_{rs} in the previous Proposition, it is easy to see that

$$\varphi_+(L_{0+}^* \mathcal{G}_{ts}^{(n)}) \neq 0 \Rightarrow s = t \text{ and } n + 2t = 0.$$

Hence the only thing left to do is to prove that L_{0+} is orthogonal to $\mathcal{G}_{tt}^{(-2t)}$ for $t \in \mathbb{Z}_0$. But suppose this were not so. Then as $\mathcal{G}^{(-2t)}$ and $\mathcal{G}^{(0)}$ are irreducible, we would necessarily get that they are isomorphic, again by the orthogonality relations. As the inner product of L_{0+} with all $\mathcal{G}_{rs}^{(-2t)}$ except $r, s = t$ is zero, this would then imply that L_{0+} must be a scalar multiple of $\mathcal{G}_{tt}^{(-2t)}$, which is equivalent with saying that $p_t(b^* b; 1, 0 \mid q^2)$ is a scalar multiple of the unit. As the spectrum of $b^* b$ is not finite, and $p_t(x; 1, 0 \mid q^2)$ is a non-constant polynomial in x , we obtain a contradiction. Hence the $\mathcal{G}^{(n)}$ are mutually non-isomorphic.

We end by showing that the $\mathcal{G}_{ts}^{(n)}$ have a σ -weakly dense linear span in N . Consider, for $k, t \in \mathbb{N}$, the element $\mathcal{G}_{t, t+k}^{(-2t-k)}$. Then, up to a non-zero constant, this equals the element $L_{0+} b^k p_t(b^* b; q^{2k}, 0 \mid q^2)$. As the $p_t(x; q^{2k}, 0 \mid q^2)$ are polynomials of degree t , we conclude that the linear span of the $\mathcal{G}_{t, t+k}^{(-2t-k)}$ contains all elements of the form $L_{0+} b^{k+t} (b^*)^t$. A similar argument shows that the $\mathcal{G}_{s+k, s}^{(-2s-k)}$ contain all elements of the form $L_{0+} b^s (b^*)^{s+k}$. Hence the linear span of all $\mathcal{G}_{ts}^{(n)}$ contains all elements of the form $L_{0+} b^s (b^*)^t$ for $s, t \in \mathbb{N}$. As this linear span is closed under left multiplication with powers of v , we conclude that this linear span contains all elements of the form $v^n L_{0+} b^s (b^*)^t$ where $n \in \mathbb{Z}$, $s, t \in \mathbb{N}$. As we can σ -weakly approximate elements of the form $e_{rr} \otimes 1 \otimes S^k$ by elements in the linear span of the $b^s (b^*)^t$, it follows immediately that the σ -weak closure of the linear span of the $\mathcal{G}_{ts}^{(n)}$ indeed equals $N = B(l^2(\mathbb{N}), l^2(\mathbb{Z})) \otimes 1 \otimes \mathcal{L}(\mathbb{Z})$. \square

Corollary 8.6. 1. Up to isomorphism, there is only one irreducible $[(N, \Delta_N)]$ -corepresentation of $\mathcal{L}^\infty(SU_q(2))$.

2. The von Neumann algebra \widehat{N} (cf. Proposition 1.10) can be identified with $\bigoplus_{r \in \mathbb{Z}} B(l^2(\mathbb{N}))$.

Proof. The first statement follows immediately from the previous proposition, Theorem 3.2.2 and Proposition 3.8, as any $\mathcal{G}^{(n)}$ clearly implements the same irreducible $[(N, \Delta_N)]$ -corepresentation. The second statement follows from Corollary 3.3. \square

Proposition 8.7. Denote $M = \mathcal{L}^\infty(SU_q(2))$. The elements $\frac{1}{q^t \sqrt{1-q^2}} \mathcal{G}_{ts}^{(n)} \xi_M$ form an orthonormal basis of $\mathcal{L}^2(N)$, and, under the identification $\mathcal{L}^2(N) \rightarrow \bigoplus_{n \in \mathbb{Z}} (l^2(\mathbb{N}) \otimes l^2(\mathbb{N}))$ by sending $\frac{1}{q^t \sqrt{1-q^2}} \mathcal{G}_{ts}^{(n)} \xi_M$ to $e_{n,t} \otimes e_{n,s}$, the element

$$\sum_{n \in \mathbb{Z}} \sum_{i,j=0}^{\infty} 1 \otimes e_{n,ts} \otimes \mathcal{G}_{ts}^{(n)} \in \widehat{N}' \bar{\otimes} N$$

equals \widetilde{V} .

Proof. Remark first that $\varphi_+((\mathcal{G}_{ts}^{(n)})^* \mathcal{G}_{ru}^{(n)}) = \varphi_+(\mathcal{G}_{ts}^* \mathcal{G}_{ru})$. But as we have that

$$\alpha(e_{ij}) = \sum_{k,l \in \mathbb{N}} \mathcal{G}_{ik}^* \mathcal{G}_{jl} \otimes e_{kl}$$

and $(\varphi_+ \otimes \iota)\alpha = \phi_\alpha$, it follows immediately that $\varphi_+(\mathcal{G}_{ik}^* \mathcal{G}_{jl}) = \delta_{kl} \delta_{ij} (1 - q^2) q^{2i}$. Combined with the previous proposition, this proves that the $\frac{1}{q^t \sqrt{1-q^2}} \mathcal{G}_{ts}^{(n)} \xi_M$ form an orthonormal basis.

Now

$$\widetilde{V} \mathcal{G}_{ij}^{(n)} \xi_M \otimes \xi_M = \sum_{k \in \mathbb{N}} \mathcal{G}_{ik}^{(n)} \xi_M \otimes \mathcal{G}_{kj}^{(n)} \xi_M.$$

On the other hand, denoting $\widetilde{V}' = \sum_{n \in \mathbb{Z}} \sum_{i,j=0}^{\infty} 1 \otimes e_{n,ts} \otimes \mathcal{G}_{ts}^{(n)}$, we have

$$\begin{aligned} \widetilde{V}' \mathcal{G}_{ij}^{(n)} \xi_M \otimes \xi_M &\cong q^i \sqrt{1-q^2} \sum_{k \in \mathbb{N}} e_{n,i} \otimes e_{n,k} \otimes \mathcal{G}_{kj}^{(n)} \xi_M \\ &\cong \sum_{k \in \mathbb{N}} \mathcal{G}_{ik}^{(n)} \xi_M \otimes \mathcal{G}_{kj}^{(n)} \xi_M. \end{aligned}$$

As ξ_M is separating for N , this shows that $\widetilde{V} = \widetilde{V}'$. □

One may deduce from this that the ordinary matrix units $e_{n,ts}$ in $\widehat{N} = \bigoplus_{n \in \mathbb{N}} B(l^2(\mathbb{N}))$ are of the form we required in the first section, namely their corresponding vectors $e_{n,t} \in l^2(\mathbb{N})$ are eigenvectors for the trace class operator T implementing ϕ_α (it should be remarked that we are implicitly using Proposition 3.11 here).

We now remark that in [9], we had already computed that the reflection of $\mathcal{L}^\infty(SU_q(2))$ across (N, Δ_N) (see section 4 for the terminology) may be identified with Woronowicz's quantum group $\mathcal{L}^\infty(\widetilde{E}_q(2))$ (see [35]), which has as its associated von Neumann algebra

$$\mathcal{L}^\infty(\widetilde{E}_q(2)) = B(l^2(\mathbb{Z})) \bar{\otimes} 1 \bar{\otimes} \mathcal{L}(\mathbb{Z}) \subseteq B(\mathcal{L}^2(N)).$$

Now it is known (see [2]) that this is a unimodular quantum group, with its invariant nsf weight φ_0 determined by

$$\varphi_0(e_{ij} \otimes S^k) = \delta_{ij} \delta_{k,0} q^{2i}.$$

Proposition 8.8. The modular element $\delta_{\widehat{N}}$ (see Theorem 2.9.2) equals $\bigoplus_{n \in \mathbb{Z}} q^{2n} T^2 \in \widehat{N}$, where $T \in B(l^2(\mathbb{N}))$ is the operator $T e_i = q^{2i} e_i$.

Proof. Denote again by A_n the n -th component of $\delta_{\widehat{N}}$ in \widehat{N} . Then by Proposition 4.3.3, we know that $A_n = d_n^2 T^2$ for some $d_n > 0$. Moreover, by Proposition 8.7 and Theorem 4.2, we know that $\varphi_0(\mathcal{G}_{00}^{(n)} (\mathcal{G}_{00}^{(n)})^*) = \frac{1}{(1-q^2) d_n^2}$. So to know d_n , we should compute $\varphi_0(v^n L_{0+} L_{0+}^* (v^*)^n)$. But as $\sigma_t^{\varphi_0}(v) = q^{-2it} v$, we have that $\varphi_0(v^n L_{0+} L_{0+}^* (v^*)^n) = q^{-2n} \varphi_0(L_{0+} L_{0+}^*)$. As the d_n are only determined up to a fixed scalar multiple anyway, we see that we may take $d_n = q^n$, which ends the proof. □

Proposition 8.9. For all $m, n \in \mathbb{Z}$ and $i, j, k, l \in \mathbb{N}$, we have

$$\varphi_0(\mathcal{G}_{ij}^{(n)}(\mathcal{G}_{kl}^{(m)})^*) = \delta_{mn}\delta_{ik}\delta_{jl}\frac{1}{q^{2n+2j}}.$$

Proof. By Proposition 4.2 and the previous proposition, we have that

$$\varphi_0(\mathcal{G}_{ij}^{(n)}(\mathcal{G}_{kl}^{(m)})^*) = \delta_{mn}\delta_{ik}\delta_{jl}\frac{c}{q^{2n+2j}}$$

for a certain constant c . This constant is precisely the number $\varphi_0(L_{0+}L_{0+}^*)$ which we neglected to compute in the previous proposition. But

$$\begin{aligned} \varphi_0(L_{0+}L_{0+}) &= \sum_{k \in \mathbb{Z}} q^{2k}(q^{2k+2}; q^2)_\infty \\ &= (q^2; q^2)_\infty \sum_{k \in \mathbb{N}} \frac{q^{2k}}{(q^2; q^2)_k} \\ &= 1, \end{aligned}$$

by the q -binomial theorem. □

Remark: These orthogonality relations can also be written out in terms of the Wall polynomials p_n . Then one would simply get back the well-known orthogonality relations between these polynomials (see e.g. [19], equation (2.6)), from which the above orthogonality relations can also be directly deduced.

As a final computation, let us determine the fusion rules between the $\mathcal{G}^{(n)}$ and the irreducible corepresentations U_r of $\mathcal{L}^\infty(SU_q(2))$ (where $r \in \frac{1}{2}\mathbb{N}$).

Proposition 8.10. For all $n \in \mathbb{Z}$ and $r \in \frac{1}{2}\mathbb{N}$, we have

$$\mathcal{G}^{(n)} \times U_r \cong \bigoplus_{i=0}^{2r} \mathcal{G}^{(n-2r+2i)}.$$

Proof. By multiplying to the left with powers of v , it is easy to see that the fusion rules are invariant under translation of n . We may therefore restrict to the case $n = 0$.

It is then well-known that if we consider $\mathcal{L}^\infty(S_{q^0}^2)$ as an $\mathcal{L}^\infty(SU_q(2))$ -comodule, the coaction α splits as $\bigoplus_{s \in \frac{1}{2}\mathbb{N}} U_{2s}$. From the proof of Proposition 3.9 and the remark preceding it, we obtain that $\mathcal{G} \times U_r$ will then split as a direct sum of less than $2r + 1$ corepresentations. By the orthogonality relations between the $\mathcal{G}_{ij}^{(n)}$, it then suffices to find in each $\mathcal{G}^{(2i-2r)}$, with $0 \leq i \leq 2r$, a component which has non-trivial scalar product with a matrix element of $\mathcal{G} \times U_r$.

By looking at the border of U_r , and by using $G_{00}^{(0)} = L_{0+}$, we find in $\mathcal{G} \times U_r$ the elements $L_{0+}a^i b^{2r-i}$ (up to a non-zero scalar), where $i \in \mathbb{N}$ with $i \leq 2r$. It is then enough to find inside $\mathcal{G}^{(2i-2r)}$ some element which has non-trivial scalar product with $L_{0+}a^i b^{2r-i}$. But by an easy computation, we have $L_{0+}a = vL_{0+}(1 - b^*b)$, and then by induction $L_{0+}a^i b^{2r-i} = v^i L_{0+}(q^{-2i}b^*b; q^2)_i b^{2r-i}$. On the other hand, $\mathcal{G}_{0,2r-i}^{(2i-2r)}$ equals $v^i L_{0+}b^{2r-i}$, up to a scalar. As $(q^{-2i}b^*b; q^2)_i (b^*b)^{2r-i} L_{0+}^* L_{0+}$ is a non-zero positive operator, we find that indeed $\varphi_+((\mathcal{G}_{0,2r-i}^{(2i-2r)})^* L_{0+}a^i b^{2r-i}) \neq 0$, which then finishes the proof. □

Remarks:

1. By the discussion following Proposition 3.9, we could also have deduced these fusion rules directly from [28], as the multiplicity diagram of the ergodic coaction of $SU_q(2)$ on the standard Podleś sphere is explicitly computed there.
2. In [11], we discussed the ‘reflection technique’ (cf. section 4) with respect to another action of $SU_q(2)$ on a type I factor, namely the von Neumann algebraic completion of its action on the ‘quantum projective plane’ (see e.g. [15]). We showed that the reflected quantum group in this case was the extended $\widetilde{SU}_q(1,1)$ quantum group of Koelink and Kustermans ([18]). This shows in particular that the Galois co-object $(\widetilde{N}, \Delta_{\widetilde{N}})$ constructed from the quantum projective plane action is different from the one we considered in this section. In fact, as the multiplicity diagram of this action was explicitly computed in [28], we see that the $(\widetilde{N}, \Delta_{\widetilde{N}})$ -projective representations of $SU_q(2)$ are labeled by the forked half-line $\{0_+, 0_-\} \cup \mathbb{N}_0$ (again by the discussion following Proposition 3.9). Now it can be shown that the quantum group $\widetilde{SU}_q(1,1)$ contains only two group-like unitaries. By Proposition 3.5 of [10], this implies that the associated $[(\widetilde{N}, \Delta_{\widetilde{N}})]$ -projective corepresentations still form a (countably) infinite family (which in fact will be parametrized by \mathbb{N} , as obtained from $\{0_+, 0_-\} \cup \mathbb{N}_0$ by identifying 0_+ with 0_-). We do not know if this family of ‘ergodic actions on type I factors’ already occurs somewhere in the literature. At the moment, we have not even succeeded in explicitly describing these ergodic coactions, except for two cases. One case is the action on the quantum projective plane itself, which we will denote as α . The other coaction $\tilde{\alpha}$ is obtained by amplifying with the spin $1/2$ -representation $U_{1/2}$ of $SU_q(2)$:

$$\tilde{\alpha} : B(\mathcal{H}_\alpha) \otimes M_2(\mathbb{C}) \rightarrow M \bar{\otimes} B(\mathcal{H}_\alpha) \otimes M_2(\mathbb{C}) : x \rightarrow (U_{1/2})_{13}^*(\alpha \otimes \iota)(x)(U_{1/2})_{13}.$$

Note that this is still irreducible, as the spectral decomposition of α only contains corepresentations corresponding to even integer spin by [28] (see the proof of Proposition 3.9). Remark that the ergodic coaction $\tilde{\alpha}$ is not isomorphic to a coideal of $SU_q(2)$ (again by [28]).

References

- [1] E. Aljadeff, P. Etingof, S. Gelaki and D. Nikshych, On twisting of finite-dimensional Hopf algebras, *Journal of Algebra* **256** (2) (2002), 484–501.
- [2] S. Baaj, Représentation régulière du groupe quantique des déplacements de Woronowicz, *Astérisque*, **232** (1995), 11–49.
- [3] S. Baaj and G. Skandalis, Unitaires multiplicatifs et dualité pour les produits croisés de C^* -algèbres, *Ann. Sci. École Norm. Sup., 4^e série*, **26** (4) (1993), 425–488.
- [4] J. Bichon, A. De Rijdt and S. Vaes, Ergodic coactions with large quantum multiplicity and monoidal equivalence of quantum groups, *Comm. Math. Phys.* **262** (2006), 703–728.
- [5] F. Boca, Ergodic actions of compact matrix pseudogroups on C^* -algebras, In: *Recent advances in operator algebras (Orléans, 1992)*, *Astérisque* **232**, 93–109 (1995).
- [6] S. Caenepeel, Brauer groups, Hopf algebras and Galois theory, *K-Monographs in Mathematics* **4**, Kluwer Academic Publishers, Dordrecht (1998), 488+xvi p.
- [7] K. De Commer, Galois objects for algebraic quantum groups, *Journal of Algebra* **321** (6) (2009), 1746–1785.
- [8] K. De Commer, Galois objects and the twisting of locally compact quantum groups, *to appear in the Journal of Operator theory*, preprint available at arXiv:math.OA/0804.2405v3.

- [9] K. De Commer, On a Morita equivalence between the duals of quantum $SU(2)$ and quantum $\tilde{E}(2)$, arXiv:math.QA/0912.4350v2.
- [10] K. De Commer, Comonoidal W^* -Morita equivalence for von Neumann bialgebras, arXiv:math.OA/1004.0824v1.
- [11] K. De Commer, On a correspondence between quantum $SU(2)$, quantum $\tilde{E}(2)$ and extended quantum $SU(1,1)$, arXiv:math.QA/1004.4307v1.
- [12] M. Enock and L. Vainerman, Deformation of a Kac algebra by an Abelian subgroup, *Comm. Math. Phys.* **178** (1996), 571–596.
- [13] P. Etingof and S. Gelaki, The classification of triangular semisimple and cosemisimple Hopf algebras over an algebraically closed field, *Internat. Math. Res. Notices* **5** (2000), 223–234.
- [14] G. Gasper and M. Rahman, Basic Hypergeometric Series, *Cambridge University Press*, Cambridge, U.K. (1990).
- [15] P. Hajac, R. Matthes, W. Szymański, Quantum Real Projective Space, Disc and Spheres, *Algebras and Representation Theory* **6** (2) (2003), 169–192.
- [16] A. Ioana, S. Popa and S. Vaes, A class of superrigid group von Neumann algebras, *in preparation*.
- [17] M. Izumi and H. Kosaki, On a subfactor analogue of the second cohomology, *Rev. Math. Phys.* **14** (2002), 733–757.
- [18] E. Koelink and J. Kustermans, A locally compact quantum group analogue of the normalizer of $SU(1,1)$ in $SL(2, \mathbb{C})$, *Comm. Math. Phys.* **233** (2003), 231–296.
- [19] T. H. Koornwinder, The addition formula for little q -Legendre polynomials and the $SU(2)$ quantum group, *SIAM J. Math. Anal.* **22** (1) (1991), 295–301.
- [20] J. Kustermans and S. Vaes, Locally compact quantum groups in the von Neumann algebraic setting, *Mathematica Scandinavica* **92** (1) (2003), 68–92.
- [21] M. Landstad, Operator algebras and compact groups, in *Operator algebras and Group Representations*, Vol. II, Monographs Stud. Math. **18**, Pitman (1984), 33–47.
- [22] J. Lining, Representation and duality of unimodular C^* -discrete quantum groups, *J. Korean Math. Soc.* **45** (2) (2008), 575–585.
- [23] T. Masuda and Y. Nakagami, A von Neumann algebra framework for the duality of the quantum groups, *Publ. RIMS, Kyoto University* **30** (1994), 799–850.
- [24] M. Movshev, Twisting in group algebras of finite groups (Russian), *Funktsional. Anal. i Prilozhen* **27** (1993), 17–23, English translation in *Funct. Anal. Appl.* **27** (1993), 240–244.
- [25] J. Quigg, Discrete C^* -coactions and C^* -algebraic bundles, *J. Austral. Math. Soc. Ser. A* **60** (2) (1996), 204–221.
- [26] P. Schauenburg, Hopf-Galois and Bi-Galois extensions, *Galois theory, Hopf algebras, and semi-abelian categories*, *Fields Inst. Commun.* **43**, AMS (2004), 469–515.
- [27] M. Takesaki, Theory of Operator Algebras II, *Springer, Berlin* (2003).
- [28] R. Tomatsu, Compact quantum ergodic systems, *J. Funct. Anal.* **254** (2008), 1–83.
- [29] S. Vaes, The unitary implementation of a locally compact quantum group action, *Journal of Functional Analysis*. **180** (2001), 426–480.
- [30] S. Vaes and N. Vander Vennet, Identification of the Poisson and Martin boundaries of orthogonal discrete quantum groups, *Journal of the Institute of Mathematics of Jussieu* **7** (2008), 391–412.
- [31] L. Vainerman, 2-cocycles and twisting of Kac algebras, *Comm. Math. Phys.* **191** (1998), 697–721.

- [32] A. Van Daele, Morita equivalence of algebraic quantum groups, *in preparation*.
- [33] A. Wassermann, Ergodic actions of compact groups on operator algebras. II. Classification of full multiplicity ergodic actions, *Canad. J. Math.* **40** (1988), 1482-1527.
- [34] S.L. Woronowicz, Compact matrix pseudogroups, *Comm. Math. Phys.* **111** (1987), 613–665.
- [35] S.L. Woronowicz, Quantum $E(2)$ -group and its Pontryagin dual, *Letters on Math. Phys.* **23** (1991), 251–263.
- [36] S.L. Woronowicz, Compact quantum groups, *in: Symétries quantiques (Les Houches, 1995)*, North-Holland (1998), 845-884.